Duality of Linear Programming

René Thiemann

March 17, 2025

Abstract

We formalize the weak and strong duality theorems of linear programming. For the strong duality theorem we provide three sufficient preconditions: both the primal problem and the dual problem are satisfiable, the primal problem is satisfiable and bounded, or the dual problem is satisfiable and bounded. The proofs are based on an existing formalization of Farkas' Lemma.

Contents

1	Introduction	1
2	Minimum and Maximum of Potentially Infinite Sets	2
3	Weak and Strong Duality of Linear Programming	3

1 Introduction

The proofs are taken from a textbook on linear programming [3]. There clearly is already an related AFP entry on linear programming [2] and we briefly explain the relationship between that entry and this one.

- The other AFP entry provides an algorithm for solving linear programs based on an existing simplex implementation. Since the simplex implementation is formulated only for rational numbers, several results are only available for rational numbers. Moreover, the simplex algorithm internally works on sets of inequalities that are represented by linear polynomials, and there are conversions between matrix-vector inequalities and linear polynomial inequalities. Finally, that AFP entry does not contain the strong duality theorem, which is the essential result in this AFP entry.
- This AFP entry has completely been formalized in the matrix-vector representation. It mainly consists of the strong duality theorems without any algorithms. The proof of these theorems are based on Farkas'

Lemma which is provided in [1] for arbitrary linearly ordered fields. Therefore, also the duality theorems are proven in that generality without the restriction to rational numbers.

2 Minimum and Maximum of Potentially Infinite Sets

theory Minimum-Maximum imports Main begin

We define minima and maxima of sets. In contrast to the existing *Min* and *Max* operators, these operators are not restricted to finite sets

definition Maximum :: 'a :: linorder set \Rightarrow 'a where Maximum $S = (THE x. x \in S \land (\forall y \in S. y \leq x))$ **definition** Minimum :: 'a :: linorder set \Rightarrow 'a where Minimum $S = (THE x. x \in S \land (\forall y \in S. x \leq y))$

definition has-Maximum where has-Maximum $S = (\exists x. x \in S \land (\forall y \in S. y \leq x))$ **definition** has-Minimum where has-Minimum $S = (\exists x. x \in S \land (\forall y \in S. x \leq x))$

definition has-Minimum where has-Minimum $S = (\exists x. x \in S \land (\forall y \in S. x \leq y))$

```
lemma eqMaximumI:

assumes x \in S

and \bigwedge y. y \in S \implies y \leq x

shows Maximum S = x

\langle proof \rangle

lemma eqMinimumI:

assumes x \in S
```

and $\bigwedge y. y \in S \Longrightarrow x \leq y$ shows Minimum S = x $\langle proof \rangle$

lemma has-MaximumD: assumes has-Maximum S shows Maximum $S \in S$ $x \in S \Longrightarrow x \leq Maximum S$ $\langle proof \rangle$

```
lemma has-MinimumD:

assumes has-Minimum S

shows Minimum S \in S

x \in S \Longrightarrow Minimum S \leq x

\langle proof \rangle
```

On non-empty finite sets, Minimum and Min coincide, and similarly Maxi-

mum and Max.

lemma Minimum-Min: **assumes** finite $S S \neq \{\}$ **shows** Minimum S = Min S $\langle proof \rangle$ **lemma** Maximum-Max: **assumes** finite $S S \neq \{\}$

shows Maximum S = Max S $\langle proof \rangle$

For natural numbers, having a maximum is the same as being bounded from above and non-empty, or being finite and non-empty.

lemma has-Maximum-nat-iff-bdd-above: has-Maximum (A :: nat set) \longleftrightarrow bdd-above $A \land A \neq \{\} \ \langle proof \rangle$

lemma has-Maximum-nat-iff-finite: has-Maximum (A :: nat set) \longleftrightarrow finite $A \land A \neq \{\}$

 $\langle proof \rangle$

lemma bdd-above-Maximum-nat: $(x :: nat) \in A \Longrightarrow bdd$ -above $A \Longrightarrow x \le Maximum A \land proof \rangle$

end

3 Weak and Strong Duality of Linear Programming

theory LP-Duality imports Linear-Inequalities.Farkas-Lemma Minimum-Maximum begin

lemma *weak-duality-theorem*:

```
fixes A :: 'a :: linordered-comm-semiring-strict mat

assumes A: A \in carrier-mat nr nc

and b: b \in carrier-vec nr

and c: c \in carrier-vec nc

and x: x \in carrier-vec nc

and Axb: A *_v x \leq b

and y0: y \geq 0_v nr

and yA: A^T *_v y = c

shows c \cdot x \leq b \cdot y

\langle proof \rangle
```

corollary *unbounded-primal-solutions*: **fixes** A :: 'a :: *linordered-idom mat*

```
assumes A: A \in carrier-mat \ nr \ nc
and b: b \in carrier-vec \ nr
and c: c \in carrier-vec \ nc
and unbounded: \forall v. \exists x \in carrier-vec \ nc. \ A *_v x \leq b \land c \cdot x \geq v
shows \neg (\exists y. y \geq 0_v \ nr \land A^T *_v y = c)
\langle proof \rangle
```

corollary unbounded-dual-solutions: **fixes** A :: 'a :: linordered-idom mat **assumes** $A: A \in carrier-mat$ nr nc **and** $b: b \in carrier-vec$ nr **and** $c: c \in carrier-vec$ nc **and** unbounded: $\forall v. \exists y. y \ge 0_v nr \land A^T *_v y = c \land b \cdot y \le v$ **shows** $\neg (\exists x \in carrier-vec nc. A *_v x \le b)$ $\langle proof \rangle$

A version of the strong duality theorem which demands that both primal and dual problem are solvable. At this point we do not use min- or maxoperations

theorem strong-duality-theorem-both-sat: **fixes** A :: 'a :: trivial-conjugatable-linordered-field mat **assumes** $A: A \in carrier-mat$ nr nc **and** $b: b \in carrier-vec$ nr **and** $c: c \in carrier-vec$ nc **and** $primal: \exists x \in carrier-vec$ nc. $A *_v x \leq b$ **and** $dual: \exists y. y \geq 0_v nr \wedge A^T *_v y = c$ **shows** $\exists x y$. $x \in carrier-vec$ $nc \wedge A *_v x \leq b \wedge$ $y \geq 0_v nr \wedge A^T *_v y = c \wedge$ $c \cdot x = b \cdot y$ $\langle proof \rangle$

A version of the strong duality theorem which demands that the primal problem is solvable and the objective function is bounded.

```
theorem strong-duality-theorem-primal-sat-bounded:

fixes bound :: 'a :: trivial-conjugatable-linordered-field

assumes A: A \in carrier-mat nr nc

and b: b \in carrier-vec nr

and c: c \in carrier-vec nc.

and sat: \exists x \in carrier-vec nc. A *_v x \leq b

and bounded: \forall x \in carrier-vec nc. A *_v x \leq b \longrightarrow c \cdot x \leq bound

shows \exists x y.

x \in carrier-vec nc \land A *_v x \leq b \land

y \geq 0_v nr \land A^T *_v y = c \land

c \cdot x = b \cdot y

\langle proof \rangle
```

A version of the strong duality theorem which demands that the dual problem is solvable and the objective function is bounded. **theorem** strong-duality-theorem-dual-sat-bounded: **fixes** bound :: 'a :: trivial-conjugatable-linordered-field **assumes** A: $A \in carrier-mat$ nr nc **and** b: $b \in carrier-vec$ nr **and** c: $c \in carrier-vec$ nc **and** sat: $\exists y. y \ge 0_v$ nr $\land A^T *_v y = c$ **and** bounded: $\forall y. y \ge 0_v$ nr $\land A^T *_v y = c \longrightarrow$ bound $\le b \cdot y$ **shows** $\exists xy$. $x \in carrier-vec$ nc $\land A *_v x \le b \land$ $y \ge 0_v$ nr $\land A^T *_v y = c \land$ $c \cdot x = b \cdot y$ $\langle proof \rangle$

Now the previous three duality theorems are formulated via min/max.

```
corollary strong-duality-theorem-min-max:

fixes A :: 'a :: trivial-conjugatable-linordered-field mat

assumes A: A \in carrier-mat nr nc

and b: b \in carrier-vec nr

and c: c \in carrier-vec nc

and primal: \exists x \in carrier-vec nc. A *_v x \leq b

and dual: \exists y. y \geq 0_v nr \land A^T *_v y = c

shows Maximum \{c \cdot x \mid x. x \in carrier-vec nc \land A *_v x \leq b\}

= Minimum \{b \cdot y \mid y. y \geq 0_v nr \land A^T *_v y = c\}

and has-Maximum \{c \cdot x \mid x. x \in carrier-vec nc \land A *_v x \leq b\}

and has-Minimum \{b \cdot y \mid y. y \geq 0_v nr \land A^T *_v y = c\}

and has-Minimum \{b \cdot y \mid y. y \geq 0_v nr \land A^T *_v y = c\}

and has-Minimum \{b \cdot y \mid y. y \geq 0_v nr \land A^T *_v y = c\}
```

corollary strong-duality-theorem-primal-sat-bounded-min-max: **fixes** bound :: 'a :: trivial-conjugatable-linordered-field **assumes** A: $A \in carrier$ -mat nr nc **and** b: $b \in carrier$ -vec nr **and** c: $c \in carrier$ -vec nc. **and** sat: $\exists x \in carrier$ -vec nc. $A *_v x \leq b$ **and** bounded: $\forall x \in carrier$ -vec nc. $A *_v x \leq b \longrightarrow c \cdot x \leq b$ ound **shows** Maximum $\{c \cdot x \mid x. x \in carrier$ -vec $nc \wedge A *_v x \leq b\}$ $= Minimum \{b \cdot y \mid y. y \geq 0_v nr \wedge A^T *_v y = c\}$ **and** has-Maximum $\{c \cdot x \mid x. x \in carrier$ -vec $nc \wedge A *_v x \leq b\}$ **and** has-Minimum $\{b \cdot y \mid y. y \geq 0_v nr \wedge A^T *_v y = c\}$ **and** has-Minimum $\{b \cdot y \mid y. y \geq 0_v nr \wedge A^T *_v y = c\}$ **and** has-Minimum $\{b \cdot y \mid y. y \geq 0_v nr \wedge A^T *_v y = c\}$

corollary strong-duality-theorem-dual-sat-bounded-min-max: **fixes** bound :: 'a :: trivial-conjugatable-linordered-field **assumes** A: $A \in carrier$ -mat nr nc **and** b: $b \in carrier$ -vec nr **and** c: $c \in carrier$ -vec nc **and** sat: $\exists y. y \ge 0_v nr \land A^T *_v y = c$ **and** bounded: $\forall y. y \ge 0_v nr \land A^T *_v y = c \longrightarrow bound \le b \cdot y$ **shows** Maximum { $c \cdot x \mid x. x \in carrier$ -vec $nc \land A *_v x \le b$ } $= Minimum \{b \cdot y \mid y. y \ge 0_v nr \land A^T *_v y = c\}$

```
and has-Maximum {c \cdot x \mid x. x \in carrier-vec nc \wedge A *_v x \leq b}
and has-Minimum {b \cdot y \mid y. y \geq 0_v nr \wedge A^T *_v y = c}
\langle proof \rangle
```

 \mathbf{end}

References

- [1] R. Bottesch, A. Reynaud, and R. Thiemann. Linear inequalities. Archive of Formal Proofs, June 2019. https://isa-afp.org/entries/ Linear_Inequalities.html, Formal proof development.
- J. Parsert and C. Kaliszyk. Linear programming. Archive of Formal Proofs, Aug. 2019. https://isa-afp.org/entries/Linear_Programming. html, Formal proof development.
- [3] A. Schrijver. Theory of linear and integer programming. John Wiley & Sons, 1998.