

# Binary Multirelations

Hitoshi Furusawa and Georg Struth

March 17, 2025

## Abstract

Binary multirelations associate elements of a set with its subsets; hence they are binary relations of type  $A \times 2^A$ . Applications include alternating automata, models and logics for games, program semantics with dual demonic and angelic nondeterministic choices and concurrent dynamic logics. This proof document supports an arXiv article that formalises the basic algebra of multirelations and proposes axiom systems for them, ranging from weak bi-monoids to weak bi-quantales.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>C-Algebras</b>	<b>2</b>
2.1	C-Monoids . . . . .	2
2.2	C-Trioids . . . . .	5
2.3	Results for Concurrent Dynamic Algebra . . . . .	7
2.4	C-Lattices . . . . .	9
2.5	Domain in C-Lattices . . . . .	14
2.6	Structural Properties of C-Lattices . . . . .	16
2.7	Terminal and Nonterminal Elements . . . . .	20
2.8	Powers in C-Algebras . . . . .	25
2.9	C-Kleene Algebras . . . . .	26
2.10	C-Omega Algebras . . . . .	28
2.11	C-Nabla Algebras . . . . .	29
2.12	Proto-Quantales . . . . .	30
<b>3</b>	<b>Multirelations</b>	<b>31</b>
3.1	Basic Definitions . . . . .	32
3.2	Multirelations and Proto-Dioids . . . . .	33
3.3	Simple Properties . . . . .	33
3.4	Multirelations and C-Lattices . . . . .	36
3.5	Terminal and Nonterminal Elements . . . . .	39
3.6	Multirelations, Proto-Quantales and Iteration . . . . .	40

3.7	Further Counterexamples . . . . .	41
3.8	Relationship with Up-Closed Multirelations . . . . .	42

## 1 Introduction

This proof document contains the formal proofs for an article on *Taming Multirelations* [2]. Individual cross-references to statements in [2] have been added to this document so that both can be read in parallel. The first part of this document contains algebraic axiom systems and equational proofs. Some of these proofs are presented in a human-readable style to indicate the kind of algebraic reasoning involved. The second part contains set-theoretic reasoning with concrete multirelations. Its main purpose is to justify the algebraic development and to prepare the soundness proofs of the algebraic axiomatisations with respect to the concrete multirelational model. Set-theoretic reasoning with multirelations tends to be very tedious and showing detailed proofs has not been the aim.

The algebras of multirelations proposed are based on Peleg's multirelational semantics for concurrent dynamic logic [3]. The most basic axiom systems consider multirelations under the operations of sequential and concurrent composition with two corresponding units. These are enriched by lattice operations and various fixpoints. A main source of complexity is the set-theoretic definition of sequential composition of multirelations, which is based on higher-order logic. Its use often requires the Axiom of Choice. In addition, sequential composition is not associative.

Part of this formalisation is also relevant to a previous approach to concurrent dynamic algebra by Furusawa and Struth [1]. More material on variants of multirelations, game algebras and concurrent dynamic algebras will be added in the future.

The authors are indebted to Alasdair Armstrong and Victor Gomes for help with some tricky Isabelle proofs.

## 2 C-Algebras

```
theory C-Algebras
imports Kleene-Algebra.Diodid
begin

no-notation
times (infixl <..> 70)
```

### 2.1 C-Monoids

We start with the c-monoid axioms. These can be found in Section 4 of [2].

```
class proto-monoid =
```

```

fixes s-id :: 'a ( $\langle 1_\sigma \rangle$ )
and s-prod :: 'a  $\Rightarrow$  'a  $\Rightarrow$  'a (infixl  $\leftrightarrow$  80)
assumes s-prod-idl [simp]:  $1_\sigma \cdot x = x$ 
and s-prod-idr [simp]:  $x \cdot 1_\sigma = x$ 

class proto-bi-monoid = proto-monoid +
fixes c-id :: 'a ( $\langle 1_\pi \rangle$ )
and c-prod :: 'a  $\Rightarrow$  'a  $\Rightarrow$  'a (infixl  $\langle \parallel \rangle$  80)
assumes c-prod-idl [simp]:  $1_\pi \parallel x = x$ 
and c-prod-assoc:  $(x \parallel y) \parallel z = x \parallel (y \parallel z)$ 
and c-prod-comm:  $x \parallel y = y \parallel x$ 

class c-monoid = proto-bi-monoid +
assumes c1 [simp]:  $(x \cdot 1_\pi) \parallel x = x$ 
and c2 [simp]:  $((x \cdot 1_\pi) \parallel 1_\sigma) \cdot y = (x \cdot 1_\pi) \parallel y$ 
and c3:  $(x \parallel y) \cdot 1_\pi = (x \cdot 1_\pi) \parallel (y \cdot 1_\pi)$ 
and c4:  $(x \cdot y) \cdot 1_\pi = x \cdot (y \cdot 1_\pi)$ 
and c5 [simp]:  $1_\sigma \parallel 1_\sigma = 1_\sigma$ 

```

**begin**

Next we define domain explicitly as at the beginning of Section 4 in [2] and start proving the algebraic facts from Section 4. Those involving concrete multirelations, such as Proposition 4.1, are considered in the theory file for multirelations.

**definition (in c-monoid)** d :: '*a*  $\Rightarrow$  '*a* **where**  
 $d\ x = (x \cdot 1_\pi) \parallel 1_\sigma$

**lemma** c-prod-idr [simp]:  $x \parallel 1_\pi = x$   
 $\langle proof \rangle$

We prove the retraction properties of Lemma 4.2.

**lemma** c-idem [simp]:  $1_\pi \cdot 1_\pi = 1_\pi$   
 $\langle proof \rangle$

**lemma** d-idem [simp]:  $d(d\ x) = d\ x$   
 $\langle proof \rangle$

**lemma** p-id-idem:  $(x \cdot 1_\pi) \cdot 1_\pi = x \cdot 1_\pi$   
 $\langle proof \rangle$

Lemma 4.3.

**lemma** c2-d:  $d\ x \cdot y = (x \cdot 1_\pi) \parallel y$   
 $\langle proof \rangle$

**lemma** cd-2-var:  $d(x \cdot 1_\pi) \cdot y = (x \cdot 1_\pi) \parallel y$   
 $\langle proof \rangle$

**lemma** *dc-prop1* [*simp*]:  $d(x \cdot 1_\pi) = x \cdot 1_\pi$   
*⟨proof⟩*

**lemma** *dc-prop2* [*simp*]:  $d(x \cdot 1_\pi) = d x$   
*⟨proof⟩*

**lemma** *ds-prop* [*simp*]:  $d(x \parallel 1_\sigma) = d x$   
*⟨proof⟩*

**lemma** *dc* [*simp*]:  $d(1_\pi) = 1_\sigma$   
*⟨proof⟩*

Part (5) of this Lemma has already been verified above. The next two statements verify the two algebraic properties mentioned in the proof of Proposition 4.4.

**lemma** *dc-iso* [*simp*]:  $d(d(x \cdot 1_\pi)) = d x$   
*⟨proof⟩*

**lemma** *cd-iso* [*simp*]:  $d(x \cdot 1_\pi) \cdot 1_\pi = x \cdot 1_\pi$   
*⟨proof⟩*

Proposition 4.5.

**lemma** *d-conc6*:  $d(x \parallel y) = d x \parallel d y$   
*⟨proof⟩*

**lemma** *d-conc-s-prod-ax*:  $d(x \parallel d y) = d x \cdot d y$   
*⟨proof⟩*

**lemma** *d-rest-ax* [*simp*]:  $d(x \cdot x) = x$   
*⟨proof⟩*

**lemma** *d-loc-ax* [*simp*]:  $d(x \cdot d y) = d(x \cdot y)$   
*⟨proof⟩*

**lemma** *d-exp-ax* [*simp*]:  $d(d(x \cdot y)) = d x \cdot d y$   
*⟨proof⟩*

**lemma** *d-comm-ax*:  $d(x \cdot d y) = d y \cdot d x$   
*⟨proof⟩*

**lemma** *d-s-id-prop* [*simp*]:  $d(1_\sigma) = 1_\sigma$   
*⟨proof⟩*

Next we verify the conditions of Proposition 4.6.

**lemma** *d-s-prod-closed* [*simp*]:  $d(d(x \cdot d y)) = d x \cdot d y$   
*⟨proof⟩*

**lemma** *d-p-prod-closed* [*simp*]:  $d(d(x \parallel d y)) = d x \parallel d y$

$\langle proof \rangle$

**lemma** *d-idem2* [simp]:  $d x \cdot d x = d x$   
 $\langle proof \rangle$

**lemma** *d-assoc*:  $(d x \cdot d y) \cdot d z = d x \cdot (d y \cdot d z)$   
 $\langle proof \rangle$

**lemma** *iso-1* [simp]:  $(d x \cdot 1_\pi) \parallel 1_\sigma = d x$   
 $\langle proof \rangle$

Lemma 4.7.

**lemma** *x-c-par-idem* [simp]:  $(x \cdot 1_\pi) \parallel (x \cdot 1_\pi) = x \cdot 1_\pi$   
 $\langle proof \rangle$

**lemma** *d-idem-par* [simp]:  $d x \parallel d x = d x$   
 $\langle proof \rangle$

**lemma** *d-inter-r*:  $d x \cdot (y \parallel z) = (d x \cdot y) \parallel (d x \cdot z)$   
 $\langle proof \rangle$

Now we provide the counterexamples of Lemma 4.8.

**lemma**  $(x \parallel y) \cdot d z = (x \cdot d z) \parallel (y \cdot d z)$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $(x \cdot y) \cdot d z = x \cdot (y \cdot d z)$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $1_\pi \cdot x = 1_\pi$   
**nitpick**  
 $\langle proof \rangle$

**end**

## 2.2 C-Trioids

We can now define the class of c-trioids and prove properties in this class. This covers the algebraic material of Section 5 in [2].

**class** *proto-diodid* = *join-semilattice-zero* + *proto-monoid* +  
**assumes** *s-prod-distr*:  $(x + y) \cdot z = x \cdot z + y \cdot z$   
**and** *s-prod-subdistl*:  $x \cdot y + x \cdot z \leq x \cdot (y + z)$   
**and** *s-prod-annil* [simp]:  $0 \cdot x = 0$

**begin**

**lemma** *s-prod-isol*:  $x \leq y \implies z \cdot x \leq z \cdot y$

```

⟨proof⟩

lemma s-prod-isor:  $x \leq y \implies x \cdot z \leq y \cdot z$ 
⟨proof⟩

end

class proto-trioid = proto-dioid + proto-bi-monoid +
assumes p-prod-distl:  $x \parallel (y + z) = x \parallel y + x \parallel z$ 
and p-rpd-annir [simp]:  $x \parallel 0 = 0$ 

sublocale proto-trioid ⊆ ab-semigroup-mult c-prod
⟨proof⟩

sublocale proto-trioid ⊆ dioid-one-zero (+) (||) 1π 0 (≤) (<)
⟨proof⟩

class c-trioid = proto-trioid + c-monoid +
assumes c6:  $x \cdot 1_{\pi} \leq 1_{\pi}$ 

begin

We show that every c-trioid is a c-monoid.

subclass c-monoid ⟨proof⟩

subclass proto-trioid ⟨proof⟩

lemma 1π · 0 = 1π
nitpick
⟨proof⟩

lemma zero-p-id-prop [simp]:  $(x \cdot 0) \cdot 1_{\pi} = x \cdot 0$ 
⟨proof⟩

The following facts prove and refute properties related to sequential and parallel subidentities.

lemma d-subid:  $d x = x \implies x \leq 1_{\sigma}$ 
⟨proof⟩

lemma x ≤ 1σ ⇒ d x = x
nitpick
⟨proof⟩

lemma p-id-term:  $x \cdot 1_{\pi} = x \implies x \leq 1_{\pi}$ 
⟨proof⟩

lemma x ≤ 1π ⇒ x · 1π = x
nitpick
⟨proof⟩

```

Proposition 5.1. is covered by the theory file on multirelations. We verify the remaining conditions in Proposition 5.2.

**lemma** *dlp-ax*:  $x \leq d x \cdot x$   
 $\langle proof \rangle$

**lemma** *d-add-ax*:  $d(x + y) = d x + d y$   
 $\langle proof \rangle$

**lemma** *d-sub-id-ax*:  $d x \leq 1_\sigma$   
 $\langle proof \rangle$

**lemma** *d-zero-ax* [*simp*]:  $d 0 = 0$   
 $\langle proof \rangle$

We verify the algebraic conditions in Proposition 5.3.

**lemma** *d-absorb1* [*simp*]:  $d x + (d x \cdot d y) = d x$   
 $\langle proof \rangle$

**lemma** *d-absorb2* [*simp*]:  $d x \cdot (d x + d y) = d x$   
 $\langle proof \rangle$

**lemma** *d-dist1*:  $d x \cdot (d y + d z) = d x \cdot d y + d x \cdot d z$   
 $\langle proof \rangle$

**lemma** *d-dist2*:  $d x + (d y \cdot d z) = (d x + d y) \cdot (d x + d z)$   
 $\langle proof \rangle$

**lemma** *d-add-prod-closed* [*simp*]:  $d(d x + d y) = d x + d y$   
 $\langle proof \rangle$

The following properties are not covered in the article.

**lemma** *x-zero-prop*:  $(x \cdot 0) \parallel y = d(x \cdot 0) \cdot y$   
 $\langle proof \rangle$

**lemma** *cda-add-ax*:  $d((x + y) \cdot z) = d(x \cdot z) + d(y \cdot z)$   
 $\langle proof \rangle$

**lemma** *d-x-zero*:  $d(x \cdot 0) = (x \cdot 0) \parallel 1_\sigma$   
 $\langle proof \rangle$

Lemma 5.4 is verified below because its proofs are simplified by using facts from the next subsection.

## 2.3 Results for Concurrent Dynamic Algebra

The following proofs and refutation are related to Section 6 in [2]. We do not consider those involving Kleene algebras in this section. We also do not introduce specific notation for diamond operators.

First we prove Lemma 6.1. Part (1) and (3) have already been verified above. Part (2) and (4) require additional assumptions which are present in the context of concurrent dynamic algebra [1]. We also present the counterexamples from Lemma 6.3.

**lemma**  $(x \cdot y) \cdot d z = x \cdot (y \cdot d z)$

**nitpick**

$\langle proof \rangle$

**lemma**  $d((x \cdot y) \cdot z) = d(x \cdot d(y \cdot z))$

**nitpick**

$\langle proof \rangle$

**lemma**  $cda\text{-}ax1: (x \cdot y) \cdot d z = x \cdot (y \cdot d z) \implies d((x \cdot y) \cdot z) = d(x \cdot d(y \cdot z))$

$\langle proof \rangle$

**lemma**  $d\text{-}inter: (x \parallel y) \cdot d z = (x \cdot d z) \parallel (y \cdot d z)$

**nitpick**

$\langle proof \rangle$

**lemma**  $d((x \parallel y) \cdot z) = d(x \cdot z) \cdot d(y \cdot z)$

**nitpick**

$\langle proof \rangle$

**lemma**  $cda\text{-}ax2:$

**assumes**  $(x \parallel y) \cdot d z = (x \cdot d z) \parallel (y \cdot d z)$

**shows**  $d((x \parallel y) \cdot z) = d(x \cdot z) \cdot d(y \cdot z)$

$\langle proof \rangle$

Next we present some results that do not feature in the article.

**lemma**  $(x \cdot y) \cdot 0 = x \cdot (y \cdot 0)$

**nitpick**

$\langle proof \rangle$

**lemma**  $d\text{-}x\text{-zero-prop} [\text{simp}]: d(x \cdot 0) \cdot 1_\pi = x \cdot 0$

$\langle proof \rangle$

**lemma**  $x \leq 1_\sigma \wedge y \leq 1_\sigma \implies x \cdot y = x \parallel y$

**nitpick**

$\langle proof \rangle$

**lemma**  $x \cdot (y \parallel z) \leq (x \cdot y) \parallel (x \cdot z)$

**nitpick**

$\langle proof \rangle$

**lemma**  $x \leq x \parallel x$

**nitpick**

$\langle proof \rangle$

Lemma 5.4

**lemma** *d-lb1*:  $d x \cdot d y \leq d x$   
*(proof)*

**lemma** *d-lb2*:  $d x \cdot d y \leq d y$   
*(proof)*

**lemma** *d-glb*:  $d z \leq d x \wedge d z \leq d y \implies d z \leq d x \cdot d y$   
*(proof)*

**lemma** *d-glb-iff*:  $d z \leq d x \wedge d z \leq d y \longleftrightarrow d z \leq d x \cdot d y$   
*(proof)*

**lemma** *x-zero-le-c*:  $x \cdot 0 \leq 1_\pi$   
*(proof)*

**lemma** *p-subid-lb1*:  $(x \cdot 0) \parallel (y \cdot 0) \leq x \cdot 0$   
*(proof)*

**lemma** *p-subid-lb2*:  $(x \cdot 0) \parallel (y \cdot 0) \leq y \cdot 0$   
*(proof)*

**lemma** *p-subid-idem [simp]*:  $(x \cdot 0) \parallel (x \cdot 0) = x \cdot 0$   
*(proof)*

**lemma** *p-subid-glb*:  $z \cdot 0 \leq x \cdot 0 \wedge z \cdot 0 \leq y \cdot 0 \implies z \cdot 0 \leq (x \cdot 0) \parallel (y \cdot 0)$   
*(proof)*

**lemma** *p-subid-glb-iff*:  $z \cdot 0 \leq x \cdot 0 \wedge z \cdot 0 \leq y \cdot 0 \longleftrightarrow z \cdot 0 \leq (x \cdot 0) \parallel (y \cdot 0)$   
*(proof)*

**lemma** *x-c-glb*:  $z \cdot 1_\pi \leq x \cdot 1_\pi \wedge z \cdot 1_\pi \leq y \cdot 1_\pi \implies z \cdot 1_\pi \leq (x \cdot 1_\pi) \parallel (y \cdot 1_\pi)$   
*(proof)*

**lemma** *x-c-lb1*:  $(x \cdot 1_\pi) \parallel (y \cdot 1_\pi) \leq x \cdot 1_\pi$   
*(proof)*

**lemma** *x-c-lb2*:  $(x \cdot 1_\pi) \parallel (y \cdot 1_\pi) \leq y \cdot 1_\pi$   
*(proof)*

**lemma** *x-c-glb-iff*:  $z \cdot 1_\pi \leq x \cdot 1_\pi \wedge z \cdot 1_\pi \leq y \cdot 1_\pi \longleftrightarrow z \cdot 1_\pi \leq (x \cdot 1_\pi) \parallel (y \cdot 1_\pi)$   
*(proof)*

**end**

## 2.4 C-Lattices

We can now define c-lattices and prove the results from Section 7 in [2].

```

class pbl-monoid = proto-trioid +
  fixes U :: 'a
  fixes meet :: 'a ⇒ 'a ⇒ 'a (infixl 〈⊓〉 70)
  assumes U-def:  $x \leq U$ 
  and meet-assoc:  $(x \sqcap y) \sqcap z = x \sqcap (y \sqcap z)$ 
  and meet-comm:  $x \sqcap y = y \sqcap x$ 
  and meet-idem [simp]:  $x \sqcap x = x$ 
  and absorp1:  $x \sqcap (x + y) = x$ 
  and absorp2:  $x + (x \sqcap y) = x$ 

begin

sublocale lattice (⊓) (≤) (<) (+)
  ⟨proof⟩

lemma meet-glb:  $z \leq x \wedge z \leq y \implies z \leq x \sqcap y$ 
  ⟨proof⟩

lemma meet-prop:  $z \leq x \wedge z \leq y \longleftrightarrow z \leq x \sqcap y$ 
  ⟨proof⟩

end

class pbdl-monoid = pbl-monoid +
  assumes lat-dist1:  $x + (y \sqcap z) = (x + y) \sqcap (x + z)$ 

begin

lemma lat-dist2:  $(x \sqcap y) + z = (x + z) \sqcap (y + z)$ 
  ⟨proof⟩

lemma lat-dist3:  $x \sqcap (y + z) = (x \sqcap y) + (x \sqcap z)$ 
  ⟨proof⟩

lemma lat-dist4:  $(x + y) \sqcap z = (x \sqcap z) + (y \sqcap z)$ 
  ⟨proof⟩

lemma d-equiv-prop:  $(\forall z. z + x = z + y \wedge z \sqcap x = z \sqcap y) \implies x = y$ 
  ⟨proof⟩

end

```

The symbol  $\bar{1}_\pi$  from [2] is written nc in this theory file.

```

class c-lattice = pbdl-monoid +
  fixes nc :: 'a
  assumes cl1 [simp]:  $x \cdot \bar{1}_\pi + x \cdot nc = x \cdot U$ 
  and cl2 [simp]:  $\bar{1}_\pi \sqcap (x + nc) = x \cdot 0$ 
  and cl3:  $x \cdot (y \parallel z) \leq (x \cdot y) \parallel (x \cdot z)$ 
  and cl4:  $z \parallel z \leq z \implies (x \parallel y) \cdot z = (x \cdot z) \parallel (y \cdot z)$ 

```

```

and cl5:  $x \cdot (y \cdot (z \cdot 0)) = (x \cdot y) \cdot (z \cdot 0)$ 
and cl6 [simp]:  $(x \cdot 0) \cdot z = x \cdot 0$ 
and cl7 [simp]:  $1_\sigma \parallel 1_\sigma = 1_\sigma$ 
and cl8 [simp]:  $((x \cdot 1_\pi) \parallel 1_\sigma) \cdot y = (x \cdot 1_\pi) \parallel y$ 
and cl9 [simp]:  $((x \sqcap 1_\sigma) \cdot 1_\pi) \parallel 1_\sigma = x \sqcap 1_\sigma$ 
and cl10:  $((x \sqcap nc) \cdot 1_\pi) \parallel 1_\sigma = 1_\sigma \sqcap (x \sqcap nc) \cdot nc$ 
and cl11 [simp]:  $((x \sqcap nc) \cdot 1_\pi) \parallel nc = (x \sqcap nc) \cdot nc$ 

```

**begin**

We show that every c-lattice is a c-troid (Proposition 7.1) Proposition 7.2 is again covered by the theory for multirelations.

**subclass** *c-troid*

*<proof>*

First we verify the complementation conditions after the definition of c-lattices.

**lemma** *c-nc-comp1* [simp]:  $1_\pi + nc = U$   
*<proof>*

**lemma** *c-nc-comp2* [simp]:  $1_\pi \sqcap nc = 0$   
*<proof>*

**lemma** *c-0*:  $x \sqcap 1_\pi = x \cdot 0$   
*<proof>*

Next we verify the conditions in Proposition 7.2.

**lemma** *d-s-subid*:  $d x = x \longleftrightarrow x \leq 1_\sigma$   
*<proof>*

**lemma** *term-p-subid*:  $x \cdot 1_\pi = x \longleftrightarrow x \leq 1_\pi$   
*<proof>*

**lemma** *term-p-subid-var*:  $x \cdot 0 = x \longleftrightarrow x \leq 1_\pi$   
*<proof>*

**lemma** *vec-iff*:  $d x \cdot U = x \longleftrightarrow (x \cdot 1_\pi) \parallel U = x$   
*<proof>*

**lemma** *nc-iff1*:  $x \leq nc \longleftrightarrow x \sqcap 1_\pi = 0$   
*<proof>*

**lemma** *nc-iff2*:  $x \leq nc \longleftrightarrow x \cdot 0 = 0$   
*<proof>*

The results of Lemma 7.3 are again at the multirelational level. Hence we continue with Lemma 7.4.

**lemma** *assoc-p-subid*:  $(x \cdot y) \cdot (z \cdot 1_\pi) = x \cdot (y \cdot (z \cdot 1_\pi))$

$\langle proof \rangle$

**lemma** zero-assoc3:  $(x \cdot y) \cdot 0 = x \cdot (y \cdot 0)$   
 $\langle proof \rangle$

**lemma** x-zero-interr:  $(x \cdot 0) \parallel (y \cdot 0) = (x \parallel y) \cdot 0$   
 $\langle proof \rangle$

**lemma** p-subid-interr:  $(x \cdot z \cdot 1_\pi) \parallel (y \cdot z \cdot 1_\pi) = (x \parallel y) \cdot z \cdot 1_\pi$   
 $\langle proof \rangle$

**lemma** d-interr:  $(x \cdot d z) \parallel (y \cdot d z) = (x \parallel y) \cdot d z$   
 $\langle proof \rangle$

**lemma** subidem-par:  $x \leq x \parallel x$   
 $\langle proof \rangle$

**lemma** meet-le-par:  $x \sqcap y \leq x \parallel y$   
 $\langle proof \rangle$

Next we verify Lemma 7.5 and prove some related properties.

**lemma** x-split [simp]:  $(x \sqcap nc) + (x \sqcap 1_\pi) = x$   
 $\langle proof \rangle$

**lemma** x-split-var [simp]:  $(x \sqcap nc) + (x \cdot 0) = x$   
 $\langle proof \rangle$

**lemma** s-subid-closed [simp]:  $x \sqcap nc \sqcap 1_\sigma = x \sqcap 1_\sigma$   
 $\langle proof \rangle$

**lemma** sub-id-le-nc:  $x \sqcap 1_\sigma \leq nc$   
 $\langle proof \rangle$

**lemma** s-x-c [simp]:  $1_\sigma \sqcap (x \cdot 1_\pi) = 0$   
 $\langle proof \rangle$

**lemma** s-x-zero [simp]:  $1_\sigma \sqcap (x \cdot 0) = 0$   
 $\langle proof \rangle$

**lemma** c-nc [simp]:  $(x \cdot 1_\pi) \sqcap nc = 0$   
 $\langle proof \rangle$

**lemma** zero-nc [simp]:  $(x \cdot 0) \sqcap nc = 0$   
 $\langle proof \rangle$

**lemma** nc-zero [simp]:  $(x \sqcap nc) \cdot 0 = 0$   
 $\langle proof \rangle$

Lemma 7.6.

**lemma** *c-def* [*simp*]:  $U \cdot 0 = 1_\pi$   
 $\langle proof \rangle$

**lemma** *c-x-prop* [*simp*]:  $1_\pi \cdot x = 1_\pi$   
 $\langle proof \rangle$

**lemma** *U-idem-s-prod* [*simp*]:  $U \cdot U = U$   
 $\langle proof \rangle$

**lemma** *U-idem-p-prod* [*simp*]:  $U \parallel U = U$   
 $\langle proof \rangle$

**lemma** *U-c* [*simp*]:  $U \cdot 1_\pi = 1_\pi$   
 $\langle proof \rangle$

**lemma** *s-le-nc*:  $1_\sigma \leq nc$   
 $\langle proof \rangle$

**lemma** *nc-c* [*simp*]:  $nc \cdot 1_\pi = 1_\pi$   
 $\langle proof \rangle$

**lemma** *nc-nc* [*simp*]:  $nc \cdot nc = nc$   
 $\langle proof \rangle$

**lemma** *U-nc* [*simp*]:  $U \cdot nc = U$   
 $\langle proof \rangle$

**lemma** *nc-U* [*simp*]:  $nc \cdot U = U$   
 $\langle proof \rangle$

**lemma** *nc-nc-par* [*simp*]:  $nc \parallel nc = nc$   
 $\langle proof \rangle$

**lemma** *U-nc-par* [*simp*]:  $U \parallel nc = nc$   
 $\langle proof \rangle$

We prove Lemma 7.8 and related properties.

**lemma** *x-y-split* [*simp*]:  $(x \sqcap nc) \cdot y + x \cdot 0 = x \cdot y$   
 $\langle proof \rangle$

**lemma** *x-y-prop*:  $1_\sigma \sqcap (x \sqcap nc) \cdot y = 1_\sigma \sqcap x \cdot y$   
 $\langle proof \rangle$

**lemma** *s-nc-U*:  $1_\sigma \sqcap x \cdot nc = 1_\sigma \sqcap x \cdot U$   
 $\langle proof \rangle$

**lemma** *sid-le-nc-var*:  $1_\sigma \sqcap x \leq 1_\sigma \sqcap x \parallel nc$   
 $\langle proof \rangle$

**lemma** *s-nc-par-U*:  $1_\sigma \sqcap x \parallel nc = 1_\sigma \sqcap x \parallel U$   
*(proof)*

**lemma** *x-c-nc-split*:  $(x \cdot 1_\pi) \parallel nc = (x \sqcap nc) \cdot nc + (x \cdot 0) \parallel nc$   
*(proof)*

**lemma** *x-c-U-split*:  $(x \cdot 1_\pi) \parallel U = x \cdot U + (x \cdot 0) \parallel U$   
*(proof)*

## 2.5 Domain in C-Lattices

We now prove variants of the domain axioms and verify the properties of Section 8 in [2].

**lemma** *cl9-d* [*simp*]:  $d(x \sqcap 1_\sigma) = x \sqcap 1_\sigma$   
*(proof)*

**lemma** *cl10-d*:  $d(x \sqcap nc) = 1_\sigma \sqcap (x \sqcap nc) \cdot nc$   
*(proof)*

**lemma** *cl11-d* [*simp*]:  $d(x \sqcap nc) \cdot nc = (x \sqcap nc) \cdot nc$   
*(proof)*

**lemma** *cl10-d-var1*:  $d(x \sqcap nc) = 1_\sigma \sqcap x \cdot nc$   
*(proof)*

**lemma** *cl10-d-var2*:  $d(x \sqcap nc) = 1_\sigma \sqcap (x \sqcap nc) \cdot U$   
*(proof)*

**lemma** *cl10-d-var3*:  $d(x \sqcap nc) = 1_\sigma \sqcap x \cdot U$   
*(proof)*

We verify the remaining properties of Lemma 8.1.

**lemma** *d-U* [*simp*]:  $d U = 1_\sigma$   
*(proof)*

**lemma** *d-nc* [*simp*]:  $d nc = 1_\sigma$   
*(proof)*

**lemma** *alt-d-def-nc-nc*:  $d(x \sqcap nc) = 1_\sigma \sqcap ((x \sqcap nc) \cdot 1_\pi) \parallel nc$   
*(proof)*

**lemma** *alt-d-def-nc-U*:  $d(x \sqcap nc) = 1_\sigma \sqcap ((x \sqcap nc) \cdot 1_\pi) \parallel U$   
*(proof)*

We verify the identity before Lemma 8.2 of [2] together with variants.

**lemma** *d-def-split* [*simp*]:  $d(x \sqcap nc) + d(x \cdot 0) = d x$   
*(proof)*

**lemma** *d-def-split-var* [*simp*]:  $d(x \sqcap nc) + (x \cdot 0) \parallel 1_\sigma = d x$   
 $\langle proof \rangle$

**lemma** *ax7* [*simp*]:  $(1_\sigma \sqcap x \cdot U) + (x \cdot 0) \parallel 1_\sigma = d x$   
 $\langle proof \rangle$

Lemma 8.2.

**lemma** *dom12-d*:  $d x = 1_\sigma \sqcap (x \cdot 1_\pi) \parallel nc$   
 $\langle proof \rangle$

**lemma** *dom12-d-U*:  $d x = 1_\sigma \sqcap (x \cdot 1_\pi) \parallel U$   
 $\langle proof \rangle$

**lemma** *dom-def-var*:  $d x = (x \cdot U \sqcap 1_\pi) \parallel 1_\sigma$   
 $\langle proof \rangle$

Lemma 8.3.

**lemma** *ax5-d* [*simp*]:  $d(x \sqcap nc) \cdot U = (x \sqcap nc) \cdot U$   
 $\langle proof \rangle$

**lemma** *ax5-0* [*simp*]:  $d(x \cdot 0) \cdot U = (x \cdot 0) \parallel U$   
 $\langle proof \rangle$

**lemma** *x-c-U-split2*:  $d x \cdot nc = (x \sqcap nc) \cdot nc + (x \cdot 0) \parallel nc$   
 $\langle proof \rangle$

**lemma** *x-c-U-split3*:  $d x \cdot U = (x \sqcap nc) \cdot U + (x \cdot 0) \parallel U$   
 $\langle proof \rangle$

**lemma** *x-c-U-split-d*:  $d x \cdot U = x \cdot U + (x \cdot 0) \parallel U$   
 $\langle proof \rangle$

**lemma** *x-U-prop2*:  $x \cdot nc = d(x \sqcap nc) \cdot nc + x \cdot 0$   
 $\langle proof \rangle$

**lemma** *x-U-prop3*:  $x \cdot U = d(x \sqcap nc) \cdot U + x \cdot 0$   
 $\langle proof \rangle$

**lemma** *d-x-nc* [*simp*]:  $d(x \cdot nc) = d x$   
 $\langle proof \rangle$

**lemma** *d-x-U* [*simp*]:  $d(x \cdot U) = d x$   
 $\langle proof \rangle$

The next properties of domain are important, but do not feature in [2].  
Proofs can be found in [1].

**lemma** *d-lhp1*:  $d x \leq d y \implies x \leq d y \cdot x$   
 $\langle proof \rangle$

**lemma** *d-lfp2*:  $x \leq d y \cdot x \implies d x \leq d y$   
 $\langle proof \rangle$

**lemma** *demod1*:  $d (x \cdot y) \leq d z \implies x \cdot d y \leq d z \cdot x$   
 $\langle proof \rangle$

**lemma** *demod2*:  $x \cdot d y \leq d z \cdot x \implies d (x \cdot y) \leq d z$   
 $\langle proof \rangle$

## 2.6 Structural Properties of C-Lattices

Now we consider the results from Section 9 and 10 in [2]. First we verify the conditions for Proposition 9.1.

**lemma** *d-meet-closed [simp]*:  $d (d x \sqcap d y) = d x \sqcap d y$   
 $\langle proof \rangle$

**lemma** *d-s-prod-eq-meet*:  $d x \cdot d y = d x \sqcap d y$   
 $\langle proof \rangle$

**lemma** *d-p-prod-eq-meet*:  $d x \parallel d y = d x \sqcap d y$   
 $\langle proof \rangle$

**lemma** *s-id-par-s-prod*:  $(x \sqcap 1_\sigma) \parallel (y \sqcap 1_\sigma) = (x \sqcap 1_\sigma) \cdot (y \sqcap 1_\sigma)$   
 $\langle proof \rangle$

**lemma** *s-id-par [simp]*:  $x \sqcap 1_\sigma \parallel x \sqcap 1_\sigma = x \sqcap 1_\sigma$   
 $\langle proof \rangle$

We verify the remaining conditions in Proposition 9.2.

**lemma** *p-subid-par-eq-meet*:  $(x \cdot 0) \parallel (y \cdot 0) = (x \cdot 0) \sqcap (y \cdot 0)$   
 $\langle proof \rangle$

**lemma** *p-subid-par-eq-meet-var*:  $(x \cdot 1_\pi) \parallel (y \cdot 1_\pi) = (x \cdot 1_\pi) \sqcap (y \cdot 1_\pi)$   
 $\langle proof \rangle$

**lemma** *x-zero-add-closed*:  $x \cdot 0 + y \cdot 0 = (x + y) \cdot 0$   
 $\langle proof \rangle$

**lemma** *x-zero-meet-closed*:  $(x \cdot 0) \sqcap (y \cdot 0) = (x \sqcap y) \cdot 0$   
 $\langle proof \rangle$

The following set of lemmas investigates the closure properties of vectors, including Lemma 9.3.

**lemma** *U-par-zero [simp]*:  $(0 \cdot c) \parallel U = 0$   
 $\langle proof \rangle$

**lemma** *U-par-s-id [simp]*:  $(1_\sigma \cdot 1_\pi) \parallel U = U$   
 $\langle proof \rangle$

**lemma** *U-par-p-id* [simp]:  $(1_\pi \cdot 1_\pi) \parallel U = U$   
*⟨proof⟩*

**lemma** *U-par-nc* [simp]:  $(nc \cdot 1_\pi) \parallel U = U$   
*⟨proof⟩*

**lemma** *d-add-var*:  $d x \cdot z + d y \cdot z = d (x + y) \cdot z$   
*⟨proof⟩*

**lemma** *d-interr-U*:  $(d x \cdot U) \parallel (d y \cdot U) = d (x \parallel y) \cdot U$   
*⟨proof⟩*

**lemma** *d-meet*:

**assumes**  $\bigwedge x y z. (x \sqcap y \sqcap 1_\sigma) \cdot z = (x \sqcap 1_\sigma) \cdot z \sqcap (y \sqcap 1_\sigma) \cdot z$   
**shows**  $d x \cdot z \sqcap d y \cdot z = (d x \sqcap d y) \cdot z$   
*⟨proof⟩*

Proposition 9.4

**lemma** *nc-zero-closed* [simp]:  $0 \sqcap nc = 0$   
*⟨proof⟩*

**lemma** *nc-s* [simp]:  $1_\sigma \sqcap nc = 1_\sigma$   
*⟨proof⟩*

**lemma** *nc-add-closed*:  $(x \sqcap nc) + (y \sqcap nc) = (x + y) \sqcap nc$   
*⟨proof⟩*

**lemma** *nc-meet-closed*:  $(x \sqcap nc) \sqcap (y \sqcap nc) = x \sqcap y \sqcap nc$   
*⟨proof⟩*

**lemma** *nc-scomp-closed*:  $((x \sqcap nc) \cdot (y \sqcap nc)) \leq nc$   
*⟨proof⟩*

**lemma** *nc-scomp-closed-alt* [simp]:  $((x \sqcap nc) \cdot (y \sqcap nc)) \sqcap nc = (x \sqcap nc) \cdot (y \sqcap nc)$   
*⟨proof⟩*

**lemma** *nc-eccomp-closed*:  $(x \sqcap nc) \parallel (y \sqcap nc) \leq nc$   
*⟨proof⟩*

**lemma** *nc-ccomp-closed-alt* [simp]:  $(x \parallel (y \sqcap nc)) \sqcap nc = x \parallel (y \sqcap nc)$   
*⟨proof⟩*

Lemma 9.6.

**lemma** *tarski-prod*:

**assumes**  $\bigwedge x. x \sqcap nc \neq 0 \implies nc \cdot ((x \sqcap nc) \cdot nc) = nc$   
**and**  $\bigwedge x y z. d x \cdot (y \cdot z) = (d x \cdot y) \cdot z$   
**shows**  $((x \sqcap nc) \cdot nc) \cdot ((y \sqcap nc) \cdot nc) = (\text{if } (y \sqcap nc) = 0 \text{ then } 0 \text{ else } (x \sqcap nc))$

$\cdot nc)$   
 $\langle proof \rangle$

We show the remaining conditions of Proposition 9.8.

**lemma** *nc-prod-aux* [*simp*]:  $((x \sqcap nc) \cdot nc) \cdot nc = (x \sqcap nc) \cdot nc$   
 $\langle proof \rangle$

**lemma** *nc-vec-add-closed*:  $((x \sqcap nc) \cdot nc + (y \sqcap nc) \cdot nc) \cdot nc = (x \sqcap nc) \cdot nc$   
 $+ (y \sqcap nc) \cdot nc$   
 $\langle proof \rangle$

**lemma** *nc-vec-par-closed*:  $((((x \sqcap nc) \cdot nc) \parallel ((y \sqcap nc) \cdot nc)) \cdot nc = ((x \sqcap nc) \cdot nc) \parallel ((y \sqcap nc) \cdot nc)$   
 $\langle proof \rangle$

**lemma** *nc-vec-par-is-meet*:

**assumes**  $\bigwedge x y z. (d x \sqcap d y) \cdot z = d x \cdot z \sqcap d y \cdot z$   
**shows**  $((x \sqcap nc) \cdot nc) \parallel ((y \sqcap nc) \cdot nc) = ((x \sqcap nc) \cdot nc) \sqcap ((y \sqcap nc) \cdot nc)$   
 $\langle proof \rangle$

**lemma** *nc-vec-meet-closed*:

**assumes**  $\bigwedge x y z. (d x \sqcap d y) \cdot z = d x \cdot z \sqcap d y \cdot z$   
**shows**  $((x \sqcap nc) \cdot nc) \sqcap ((y \sqcap nc) \cdot nc) \cdot nc = (x \sqcap nc) \cdot nc \sqcap (y \sqcap nc) \cdot nc$   
 $\langle proof \rangle$

**lemma** *nc-vec-seq-closed*:

**assumes**  $\bigwedge x. x \sqcap nc \neq 0 \implies nc \cdot ((x \sqcap nc) \cdot nc) = nc$   
**and**  $\bigwedge x y z. d x \cdot (y \cdot z) = (d x \cdot y) \cdot z$   
**shows**  $((x \sqcap nc) \cdot nc) \cdot ((y \sqcap nc) \cdot nc) \cdot nc = ((x \sqcap nc) \cdot nc) \cdot ((y \sqcap nc) \cdot nc)$   
 $\langle proof \rangle$

Proposition 10.1 and 10.2.

**lemma** *iso3* [*simp*]:  $d (d x \cdot U) = d x$   
 $\langle proof \rangle$

**lemma** *iso4* [*simp*]:  $d ((x \cdot 1_\pi) \parallel U) \cdot U = (x \cdot 1_\pi) \parallel U$   
 $\langle proof \rangle$

**lemma** *iso5* [*simp*]:  $((x \cdot 1_\pi) \parallel U) \cdot 1_\pi = x \cdot 1_\pi$   
 $\langle proof \rangle$

**lemma** *iso6* [*simp*]:  $((((x \cdot 1_\pi) \parallel U) \cdot 1_\pi) \parallel U = (x \cdot 1_\pi) \parallel U$   
 $\langle proof \rangle$

**lemma** *iso3-sharp* [*simp*]:  $d (d (x \sqcap nc) \cdot nc) = d (x \sqcap nc)$   
 $\langle proof \rangle$

**lemma** *iso4-sharp* [*simp*]:  $d ((x \sqcap nc) \cdot nc) \cdot nc = (x \sqcap nc) \cdot nc$   
 $\langle proof \rangle$

**lemma** *iso5-sharp [simp]*:  $((x \sqcap nc) \cdot 1_\pi) \parallel nc \cdot 1_\pi = (x \sqcap nc) \cdot 1_\pi$   
 *$\langle proof \rangle$*

**lemma** *iso6-sharp [simp]*:  $((x \sqcap nc) \cdot nc) \parallel nc = (x \sqcap nc) \cdot nc$   
 *$\langle proof \rangle$*

We verify Lemma 15.2 at this point, because it is helpful for the following proofs.

**lemma** *uc-par-meet*:  $x \parallel U \sqcap y \parallel U = x \parallel U \parallel y \parallel U$   
 *$\langle proof \rangle$*

**lemma** *uc-unc [simp]*:  $x \parallel U \parallel x \parallel U = x \parallel U$   
 *$\langle proof \rangle$*

**lemma** *uc-interr*:  $(x \parallel y) \cdot (z \parallel U) = (x \cdot (z \parallel U)) \parallel (y \cdot (z \parallel U))$   
 *$\langle proof \rangle$*

We verify the remaining cases of Proposition 10.3.

**lemma** *sc-hom-meet*:  $(d x \sqcap d y) \cdot 1_\pi = (d x) \cdot 1_\pi \sqcap (d y) \cdot 1_\pi$   
 *$\langle proof \rangle$*

**lemma** *sc-hom-seq*:  $(d x \cdot d y) \cdot 1_\pi = (d x \sqcap d y) \cdot 1_\pi$   
 *$\langle proof \rangle$*

**lemma** *cs-hom-meet*:  $d (x \cdot 1_\pi \sqcap y \cdot 1_\pi) = d (x \cdot 1_\pi) \sqcap d (y \cdot 1_\pi)$   
 *$\langle proof \rangle$*

**lemma** *sv-hom-meet*:  $(d x \sqcap d y) \cdot U = (d x) \cdot U \sqcap (d y) \cdot U$   
 *$\langle proof \rangle$*

**lemma** *sv-hom-par*:  $(x \parallel y) \cdot U = (x \cdot U) \parallel (y \cdot U)$   
 *$\langle proof \rangle$*

**lemma** *vs-hom-meet*:  $d (((x \cdot 1_\pi) \parallel U) \sqcap ((y \cdot 1_\pi) \parallel U)) = d ((x \cdot 1_\pi) \parallel U) \sqcap d ((y \cdot 1_\pi) \parallel U)$   
 *$\langle proof \rangle$*

**lemma** *cv-hom-meet*:  $(x \cdot 1_\pi \sqcap y \cdot 1_\pi) \parallel U = (x \cdot 1_\pi) \parallel U \sqcap (y \cdot 1_\pi) \parallel U$   
 *$\langle proof \rangle$*

**lemma** *cv-hom-par [simp]*:  $x \parallel U \parallel y \parallel U = (x \parallel y) \parallel U$   
 *$\langle proof \rangle$*

**lemma** *vc-hom-meet*:  $((x \cdot 1_\pi) \parallel U \sqcap (y \cdot 1_\pi) \parallel U) \cdot 1_\pi = ((x \cdot 1_\pi) \parallel U) \cdot 1_\pi$   
 $\sqcap ((y \cdot 1_\pi) \parallel U) \cdot 1_\pi$   
 *$\langle proof \rangle$*

**lemma** *vc-hom-seq*:  $((x \cdot 1_\pi) \parallel U) \cdot ((y \cdot 1_\pi) \parallel U)) \cdot 1_\pi = (((x \cdot 1_\pi) \parallel U) \cdot 1_\pi) \cdot (((y \cdot 1_\pi) \parallel U) \cdot 1_\pi)$   
 $\langle proof \rangle$

Proposition 10.4.

**lemma** *nsv-hom-meet*:  $(d x \sqcap d y) \cdot nc = (d x) \cdot nc \sqcap (d y) \cdot nc$   
 $\langle proof \rangle$

**lemma** *nsv-hom-par*:  $(x \parallel y) \cdot nc = (x \cdot nc) \parallel (y \cdot nc)$   
 $\langle proof \rangle$

**lemma** *vec-p-prod-meet*:  $((x \sqcap nc) \cdot nc) \parallel ((y \sqcap nc) \cdot nc) = ((x \sqcap nc) \cdot nc) \sqcap ((y \sqcap nc) \cdot nc)$   
 $\langle proof \rangle$

**lemma** *nvs-hom-meet*:  $d (((x \sqcap nc) \cdot nc) \sqcap ((y \sqcap nc) \cdot nc)) = d ((x \sqcap nc) \cdot nc) \sqcap d ((y \sqcap nc) \cdot nc)$   
 $\langle proof \rangle$

**lemma** *ncv-hom-meet*:  $(x \cdot 1_\pi \sqcap y \cdot 1_\pi) \parallel nc = (x \cdot 1_\pi) \parallel nc \sqcap (y \cdot 1_\pi) \parallel nc$   
 $\langle proof \rangle$

**lemma** *ncv-hom-par*:  $(x \parallel y) \parallel nc = x \parallel nc \parallel y \parallel nc$   
 $\langle proof \rangle$

**lemma** *nvc-hom-meet*:  $((x \sqcap nc) \cdot nc \sqcap (y \sqcap nc) \cdot nc) \cdot 1_\pi = ((x \sqcap nc) \cdot nc) \cdot 1_\pi \sqcap ((y \sqcap nc) \cdot nc) \cdot 1_\pi$   
 $\langle proof \rangle$

## 2.7 Terminal and Nonterminal Elements

Now we define the projection functions on terminals and nonterminal parts and verify the properties of Section 11 in [2].

**definition** *tau* ::  $'a \Rightarrow 'a (\langle \tau \rangle)$  **where**  
 $\tau x = x \cdot 0$

**definition** *nu* ::  $'a \Rightarrow 'a (\langle \nu \rangle)$  **where**  
 $\nu x = x \sqcap nc$

Lemma 11.1.

**lemma** *tau-int*:  $\tau x \leq x$   
 $\langle proof \rangle$

**lemma** *nu-int*:  $\nu x \leq x$   
 $\langle proof \rangle$

**lemma** *tau-ret* [*simp*]:  $\tau (\tau x) = \tau x$   
 $\langle proof \rangle$

**lemma** *nu-ret* [*simp*]:  $\nu (\nu x) = \nu x$   
 $\langle proof \rangle$

**lemma** *tau-iso*:  $x \leq y \implies \tau x \leq \tau y$   
 $\langle proof \rangle$

**lemma** *nu-iso*:  $x \leq y \implies \nu x \leq \nu y$   
 $\langle proof \rangle$

Lemma 11.2.

**lemma** *tau-zero* [*simp*]:  $\tau 0 = 0$   
 $\langle proof \rangle$

**lemma** *nu-zero* [*simp*]:  $\nu 0 = 0$   
 $\langle proof \rangle$

**lemma** *tau-s* [*simp*]:  $\tau 1_\sigma = 0$   
 $\langle proof \rangle$

**lemma** *nu-s* [*simp*]:  $\nu 1_\sigma = 1_\sigma$   
 $\langle proof \rangle$

**lemma** *tau-c* [*simp*]:  $\tau 1_\pi = 1_\pi$   
 $\langle proof \rangle$

**lemma** *nu-c* [*simp*]:  $\nu 1_\pi = 0$   
 $\langle proof \rangle$

**lemma** *tau-nc* [*simp*]:  $\tau nc = 0$   
 $\langle proof \rangle$

**lemma** *nu-nc* [*simp*]:  $\nu nc = nc$   
 $\langle proof \rangle$

**lemma** *tau-U* [*simp*]:  $\tau U = 1_\pi$   
 $\langle proof \rangle$

**lemma** *nu-U* [*simp*]:  $\nu U = nc$   
 $\langle proof \rangle$

Lemma 11.3.

**lemma** *tau-add* [*simp*]:  $\tau (x + y) = \tau x + \tau y$   
 $\langle proof \rangle$

**lemma** *nu-add* [*simp*]:  $\nu (x + y) = \nu x + \nu y$   
 $\langle proof \rangle$

**lemma** *tau-meet* [*simp*]:  $\tau (x \sqcap y) = \tau x \sqcap \tau y$

$\langle proof \rangle$

**lemma** *nu-meet* [simp]:  $\nu (x \sqcap y) = \nu x \sqcap \nu y$   
 $\langle proof \rangle$

**lemma** *tau-seq*:  $\tau (x \cdot y) = \tau x + \nu x \cdot \tau y$   
 $\langle proof \rangle$

**lemma** *tau-par* [simp]:  $\tau (x \parallel y) = \tau x \parallel \tau y$   
 $\langle proof \rangle$

**lemma** *nu-par-aux1*:  $x \parallel \tau y = d(\tau y) \cdot x$   
 $\langle proof \rangle$

**lemma** *nu-par-aux2* [simp]:  $\nu (\nu x \parallel \nu y) = \nu x \parallel \nu y$   
 $\langle proof \rangle$

**lemma** *nu-par-aux3* [simp]:  $\nu (\nu x \parallel \tau y) = \nu x \parallel \tau y$   
 $\langle proof \rangle$

**lemma** *nu-par-aux4* [simp]:  $\nu (\tau x \parallel \tau y) = 0$   
 $\langle proof \rangle$

**lemma** *nu-par*:  $\nu (x \parallel y) = d(\tau x) \cdot \nu y + d(\tau y) \cdot \nu x + \nu x \parallel \nu y$   
 $\langle proof \rangle$

Lemma 11.5.

**lemma** *sprod-tau-nu*:  $x \cdot y = \tau x + \nu x \cdot y$   
 $\langle proof \rangle$

**lemma** *pprod-tau-nu*:  $x \parallel y = \nu x \parallel \nu y + d(\tau x) \cdot \nu y + d(\tau y) \cdot \nu x + \tau x \parallel \tau y$   
 $\langle proof \rangle$

We now verify some additional properties which are not mentioned in the paper.

**lemma** *tau-idem* [simp]:  $\tau x \cdot \tau x = \tau x$   
 $\langle proof \rangle$

**lemma** *tau-interr*:  $(x \parallel y) \cdot \tau z = (x \cdot \tau z) \parallel (y \cdot \tau z)$   
 $\langle proof \rangle$

**lemma** *tau-le-c*:  $\tau x \leq 1_\pi$   
 $\langle proof \rangle$

**lemma** *c-le-tauc*:  $1_\pi \leq \tau 1_\pi$   
 $\langle proof \rangle$

**lemma** *x-alpha-tau* [simp]:  $\nu x + \tau x = x$

$\langle proof \rangle$

**lemma** *alpha-tau-zero* [*simp*]:  $\nu (\tau x) = 0$   
 $\langle proof \rangle$

**lemma** *tau-alpha-zero* [*simp*]:  $\tau (\nu x) = 0$   
 $\langle proof \rangle$

**lemma** *sprod-tau-nu-var* [*simp*]:  $\nu (\nu x \cdot y) = \nu (x \cdot y)$   
 $\langle proof \rangle$

**lemma** *tau-s-prod* [*simp*]:  $\tau (x \cdot y) = x \cdot \tau y$   
 $\langle proof \rangle$

**lemma** *alpha-fp*:  $\nu x = x \longleftrightarrow x \cdot 0 = 0$   
 $\langle proof \rangle$

**lemma** *alpha-prod-closed* [*simp*]:  $\nu (\nu x \cdot \nu y) = \nu x \cdot \nu y$   
 $\langle proof \rangle$

**lemma** *alpha-par-prod* [*simp*]:  $\nu (x \parallel \nu y) = x \parallel \nu y$   
 $\langle proof \rangle$

**lemma** *p-prod-tau-alpha*:  $x \parallel y = x \parallel \nu y + \nu x \parallel y + \tau x \parallel \tau y$   
 $\langle proof \rangle$

**lemma** *p-prod-tau-alpha-var*:  $x \parallel y = x \parallel \nu y + \nu x \parallel y + \tau (x \parallel y)$   
 $\langle proof \rangle$

**lemma** *alpha-par*:  $\nu (x \parallel y) = \nu x \parallel y + x \parallel \nu y$   
 $\langle proof \rangle$

**lemma** *alpha-tau* [*simp*]:  $\nu (x \cdot \tau y) = 0$   
 $\langle proof \rangle$

**lemma** *nu-par-prop*:  $\nu x = x \implies \nu (x \parallel y) = x \parallel y$   
 $\langle proof \rangle$

**lemma** *tau-seq-prop*:  $\tau x = x \implies x \cdot y = x$   
 $\langle proof \rangle$

**lemma** *tau-seq-prop2*:  $\tau y = y \implies \tau (x \cdot y) = x \cdot y$   
 $\langle proof \rangle$

**lemma** *d-nu*:  $\nu (d x \cdot y) = d x \cdot \nu y$   
 $\langle proof \rangle$

Lemma 11.6 and 11.7.

**lemma** *nu-ideal1*:  $\llbracket \nu x = x; y \leq x \rrbracket \implies \nu y = y$

$\langle proof \rangle$

**lemma** *tau-ideal1*:  $\llbracket \tau x = x; y \leq x \rrbracket \implies \tau y = y$   
 $\langle proof \rangle$

**lemma** *nu-ideal2*:  $\llbracket \nu x = x; \nu y = y \rrbracket \implies \nu(x + y) = x + y$   
 $\langle proof \rangle$

**lemma** *tau-ideal2*:  $\llbracket \tau x = x; \tau y = y \rrbracket \implies \tau(x + y) = x + y$   
 $\langle proof \rangle$

**lemma** *tau-ideal3*:  $\tau x = x \implies \tau(x \cdot y) = x \cdot y$   
 $\langle proof \rangle$

We prove the precongruence properties of Lemma 11.9.

**lemma** *tau-add-precong*:  $\tau x \leq \tau y \implies \tau(x + z) \leq \tau(y + z)$   
 $\langle proof \rangle$

**lemma** *tau-meet-precong*:  $\tau x \leq \tau y \implies \tau(x \sqcap z) \leq \tau(y \sqcap z)$   
 $\langle proof \rangle$

**lemma** *tau-par-precong*:  $\tau x \leq \tau y \implies \tau(x \parallel z) \leq \tau(y \parallel z)$   
 $\langle proof \rangle$

**lemma** *tau-seq-precongl*:  $\tau x \leq \tau y \implies \tau(z \cdot x) \leq \tau(z \cdot y)$   
 $\langle proof \rangle$

**lemma** *nu-add-precong*:  $\nu x \leq \nu y \implies \nu(x + z) \leq \nu(y + z)$   
 $\langle proof \rangle$

**lemma** *nu-meet-precong*:  $\nu x \leq \nu y \implies \nu(x \sqcap z) \leq \nu(y \sqcap z)$   
 $\langle proof \rangle$

**lemma** *nu-seq-precongr*:  $\nu x \leq \nu y \implies \nu(x \cdot z) \leq \nu(y \cdot z)$   
 $\langle proof \rangle$

We prove the congruence properties of Corollary 11.11.

**definition** *tcg* ::  $'a \Rightarrow 'a \Rightarrow \text{bool}$  **where**  
 $\text{tcg } x \ y = (\tau x \leq \tau y \wedge \tau y \leq \tau x)$

**definition** *ncg* ::  $'a \Rightarrow 'a \Rightarrow \text{bool}$  **where**  
 $\text{ncg } x \ y = (\nu x \leq \nu y \wedge \nu y \leq \nu x)$

**lemma** *tcg-refl*:  $\text{tcg } x \ x$   
 $\langle proof \rangle$

**lemma** *tcg-trans*:  $\llbracket \text{tcg } x \ y; \text{tcg } y \ z \rrbracket \implies \text{tcg } x \ z$   
 $\langle proof \rangle$

```

lemma tcg-sym:  $\text{tcg } x \ y \implies \text{tcg } y \ x$ 
   $\langle \text{proof} \rangle$ 

lemma ncg-refl:  $\text{ncg } x \ x$ 
   $\langle \text{proof} \rangle$ 

lemma ncg-trans:  $\llbracket \text{ncg } x \ y; \text{ncg } y \ z \rrbracket \implies \text{ncg } x \ z$ 
   $\langle \text{proof} \rangle$ 

lemma ncg-sym:  $\text{ncg } x \ y \implies \text{ncg } y \ x$ 
   $\langle \text{proof} \rangle$ 

lemma tcg-alt:  $\text{tcg } x \ y = (\tau \ x = \tau \ y)$ 
   $\langle \text{proof} \rangle$ 

lemma ncg-alt:  $\text{ncg } x \ y = (\nu \ x = \nu \ y)$ 
   $\langle \text{proof} \rangle$ 

lemma tcg-add:  $\tau \ x = \tau \ y \implies \tau \ (x + z) = \tau \ (y + z)$ 
   $\langle \text{proof} \rangle$ 

lemma tcg-meet:  $\tau \ x = \tau \ y \implies \tau \ (x \sqcap z) = \tau \ (y \sqcap z)$ 
   $\langle \text{proof} \rangle$ 

lemma tcg-par:  $\tau \ x = \tau \ y \implies \tau \ (x \parallel z) = \tau \ (y \parallel z)$ 
   $\langle \text{proof} \rangle$ 

lemma tcg-seql:  $\tau \ x = \tau \ y \implies \tau \ (z \cdot x) = \tau \ (z \cdot y)$ 
   $\langle \text{proof} \rangle$ 

lemma ncg-add:  $\nu \ x = \nu \ y \implies \nu \ (x + z) = \nu \ (y + z)$ 
   $\langle \text{proof} \rangle$ 

lemma ncg-meet:  $\nu \ x = \nu \ y \implies \nu \ (x \sqcap z) = \nu \ (y \sqcap z)$ 
   $\langle \text{proof} \rangle$ 

lemma ncg-seqr:  $\nu \ x = \nu \ y \implies \nu \ (x \cdot z) = \nu \ (y \cdot z)$ 
   $\langle \text{proof} \rangle$ 

end

```

## 2.8 Powers in C-Algebras

We define the power functions from Section 6 in [2] after Lemma 12.4.

```

context proto-diodid
begin

```

```

primrec p-power ::  $'a \Rightarrow \text{nat} \Rightarrow 'a$  where
   $p\text{-power } x \ 0 = 1_\sigma \mid$ 

```

```
p-power x (Suc n) = x · p-power x n
```

```
primrec power-rd :: 'a ⇒ nat ⇒ 'a where
  power-rd x 0      = 0 |
  power-rd x (Suc n) = 1σ + x · power-rd x n
```

```
primrec power-sq :: 'a ⇒ nat ⇒ 'a where
  power-sq x 0      = 1σ |
  power-sq x (Suc n) = 1σ + x · power-sq x n
```

Lemma 12.5

```
lemma power-rd-chain: power-rd x n ≤ power-rd x (n + 1)
  ⟨proof⟩
```

```
lemma power-sq-chain: power-sq x n ≤ power-sq x (n + 1)
  ⟨proof⟩
```

```
lemma pow-chain: p-power (1σ + x) n ≤ p-power (1σ + x) (n + 1)
  ⟨proof⟩
```

```
lemma pow-prop: p-power (1σ + x) (n + 1) = 1σ + x · p-power (1σ + x) n
  ⟨proof⟩
```

Next we verify facts from the proofs of Lemma 12.6.

```
lemma power-rd-le-sq: power-rd x n ≤ power-sq x n
  ⟨proof⟩
```

```
lemma power-sq-le-rd: power-sq x n ≤ power-rd x (Suc n)
  ⟨proof⟩
```

```
lemma power-sq-power: power-sq x n = p-power (1σ + x) n
  ⟨proof⟩
```

end

## 2.9 C-Kleene Algebras

The definition of c-Kleene algebra is slightly different from that in Section 6 of [2]. It is used to prove properties from Section 6 and Section 12.

```
class c-kleene-algebra = c-lattice + star-op +
  assumes star-unfold: 1σ + x · x* ≤ x*
  and star-induct: 1σ + x · y ≤ y ⇒ x* ≤ y
```

begin

```
lemma star-irr: 1σ ≤ x*
  ⟨proof⟩
```

**lemma** *star-unfold-part*:  $x \cdot x^* \leq x^*$   
*(proof)*

**lemma** *star-ext-aux*:  $x \leq x \cdot x^*$   
*(proof)*

**lemma** *star-ext*:  $x \leq x^*$   
*(proof)*

**lemma** *star-co-trans*:  $x^* \leq x^* \cdot x^*$   
*(proof)*

**lemma** *star-iso*:  $x \leq y \implies x^* \leq y^*$   
*(proof)*

**lemma** *star-unfold-eq [simp]*:  $1_\sigma + x \cdot x^* = x^*$   
*(proof)*

Lemma 12.2.

**lemma** *nu-star1*:  
**assumes**  $\bigwedge x y z. x \cdot (y \cdot z) = (x \cdot y) \cdot z$   
**shows**  $x^* \leq (\nu x)^* \cdot (1_\sigma + \tau x)$   
*(proof)*

**lemma** *nu-star2*:  
**assumes**  $\bigwedge x. x^* \cdot x^* \leq x^*$   
**shows**  $(\nu x)^* \cdot (1_\sigma + \tau x) \leq x^*$   
*(proof)*

**lemma** *nu-star*:  
**assumes**  $\bigwedge x. x^* \cdot x^* \leq x^*$   
**and**  $\bigwedge x y z. x \cdot (y \cdot z) = (x \cdot y) \cdot z$   
**shows**  $(\nu x)^* \cdot (1_\sigma + \tau x) = x^*$   
*(proof)*

Lemma 12.3.

**lemma** *tau-star*:  $(\tau x)^* = 1_\sigma + \tau x$   
*(proof)*

**lemma** *tau-star-var*:  
**assumes**  $\bigwedge x y z. x \cdot (y \cdot z) = (x \cdot y) \cdot z$   
**and**  $\bigwedge x. x^* \cdot x^* \leq x^*$   
**shows**  $\tau(x^*) = (\nu x)^* \cdot \tau x$   
*(proof)*

**lemma** *nu-star-sub*:  $(\nu x)^* \leq \nu(x^*)$   
*(proof)*

**lemma** *nu-star-nu [simp]*:  $\nu((\nu x)^*) = (\nu x)^*$

$\langle proof \rangle$

**lemma** *nu-star-tau* [*simp*]:  $\nu ((\tau x)^\star) = 1_\sigma$   
 $\langle proof \rangle$

**lemma** *tau-star-tau* [*simp*]:  $\tau ((\tau x)^\star) = \tau x$   
 $\langle proof \rangle$

**lemma** *tau-star-nu* [*simp*]:  $\tau ((\nu x)^\star) = 0$   
 $\langle proof \rangle$

Finally we verify Lemma 6.2. Proofs can be found in [1].

**lemma** *d-star-unfold* [*simp*]:  
**assumes**  $\bigwedge x y z. (x \cdot y) \cdot d z = x \cdot (y \cdot d z)$   
**shows**  $d y + d (x \cdot d (x^\star \cdot y)) = d (x^\star \cdot y)$   
 $\langle proof \rangle$

**lemma** *d-star-sim1*:  
**assumes**  $\bigwedge x y z. d z + x \cdot y \leq y \implies x^\star \cdot d z \leq y$   
**and**  $\bigwedge x y z. (x \cdot d y) \cdot z = x \cdot (d y \cdot z)$   
**and**  $\bigwedge x y z. (d x \cdot y) \cdot z = d x \cdot (y \cdot z)$   
**shows**  $x \cdot d z \leq d z \cdot y \implies x^\star \cdot d z \leq d z \cdot y^\star$   
 $\langle proof \rangle$

**lemma** *d-star-induct*:  
**assumes**  $\bigwedge x y z. d z + x \cdot y \leq y \implies x^\star \cdot d z \leq y$   
**and**  $\bigwedge x y z. (x \cdot d y) \cdot z = x \cdot (d y \cdot z)$   
**and**  $\bigwedge x y z. (d x \cdot y) \cdot z = d x \cdot (y \cdot z)$   
**shows**  $d (x \cdot y) \leq d y \implies d (x^\star \cdot y) \leq d y$   
 $\langle proof \rangle$

**end**

## 2.10 C-Omega Algebras

These structures do not feature in [2], but in fact, many lemmas from Section 13 can be proved in this setting. The proto-quantales and c-quantales using in [2] provide a more expressive setting in which least and greatest fixpoints need not be postulated; they exists due to properties of sequential composition and addition over complete lattices.

**class** *c-omega-algebra* = *c-kleene-algebra* + *omega-op* +  
**assumes** *om-unfold*:  $x^\omega \leq x \cdot x^\omega$   
**and** *om-coinduct*:  $y \leq x \cdot y \implies y \leq x^\omega$

**begin**

Lemma 13.4.

**lemma** *om-unfold-eq* [*simp*]:  $x \cdot x^\omega = x^\omega$

$\langle proof \rangle$

**lemma** *om-iso*:  $x \leq y \implies x^\omega \leq y^\omega$   
 $\langle proof \rangle$

Lemma 13.5.

**lemma** *zero-om* [*simp*]:  $0^\omega = 0$   
 $\langle proof \rangle$

**lemma** *s-id-om* [*simp*]:  $1_{\sigma^\omega} = U$   
 $\langle proof \rangle$

**lemma** *p-id-om* [*simp*]:  $1_{\pi^\omega} = 1_\pi$   
 $\langle proof \rangle$

**lemma** *nc-om* [*simp*]:  $nc^\omega = U$   
 $\langle proof \rangle$

**lemma** *U-om* [*simp*]:  $U^\omega = U$   
 $\langle proof \rangle$

Lemma 13.6.

**lemma** *tau-om1*:  $\tau x \leq \tau (x^\omega)$   
 $\langle proof \rangle$

**lemma** *tau-om2* [*simp*]:  $\tau x^\omega = \tau x$   
 $\langle proof \rangle$

**lemma** *tau-om3*:  $(\tau x)^\omega \leq \tau (x^\omega)$   
 $\langle proof \rangle$

Lemma 13.7.

**lemma** *om-nu-tau*:  $(\nu x)^\omega + (\nu x)^* \cdot \tau x \leq x^\omega$   
 $\langle proof \rangle$

**end**

## 2.11 C-Nabla Algebras

Nabla-algebras provide yet another way of formalising non-terminating behaviour in Section 13.

```
class c-nabla-algebra = c-omega-algebra +
  fixes nabla :: 'a ⇒ 'a (⟨∇⟩)
  assumes nabla-unfold: ∇ x ≤ d (x · ∇ x)
  and nabla-coinduct: d y ≤ d (x · y) ⇒ d y ≤ ∇ x
```

**begin**

**lemma** *nabla-unfold-eq* [*simp*]:  $\nabla x = d(x \cdot \nabla x)$   
*(proof)*

**lemma** *nabla-le-s*:  $\nabla x \leq 1_\sigma$   
*(proof)*

**lemma** *nabla-nu* [*simp*]:  $\nu(\nabla x) = \nabla x$   
*(proof)*

Proposition 13.9.

**lemma** *nabla-omega-U*:  
**assumes**  $\bigwedge x y z. x \cdot (d y \cdot z) = (x \cdot d y) \cdot z$   
**shows**  $(\nu x)^\omega = \nabla(\nu x) \cdot U$   
*(proof)*

Corollary 13.10.

**lemma** *nabla-omega-U-cor*:  
**assumes**  $\bigwedge x y z. x \cdot (d y \cdot z) = (x \cdot d y) \cdot z$   
**shows**  $\nabla(\nu x) \cdot U + (\nu x)^* \cdot \tau x \leq x^\omega$   
*(proof)*

Lemma 13.11.

**lemma** *nu-om-nu*:  
**assumes**  $\bigwedge x y z. x \cdot (d y \cdot z) = (x \cdot d y) \cdot z$   
**shows**  $\nu((\nu x)^\omega) = \nabla(\nu x) \cdot nc$   
*(proof)*

**lemma** *tau-om-nu*:  
**assumes**  $\bigwedge x y z. x \cdot (d y \cdot z) = (x \cdot d y) \cdot z$   
**shows**  $\tau((\nu x)^\omega) = \nabla(\nu x) \cdot 1_\pi$   
*(proof)*

Proposition 13.12.

**lemma** *wf-eq-defl*:  $(\forall y. d y \leq d(x \cdot y) \rightarrow d y = 0) \leftrightarrow (\forall y. y \leq x \cdot y \rightarrow y = 0)$   
*(proof)*

**lemma** *defl-eq-om-trivial*:  $x^\omega = 0 \leftrightarrow (\forall y. y \leq x \cdot y \rightarrow y = 0)$   
*(proof)*

**lemma** *wf-eq-om-trivial*:  $x^\omega = 0 \leftrightarrow (\forall y. d y \leq d(x \cdot y) \rightarrow d y = 0)$   
*(proof)*

**end**

## 2.12 Proto-Quantales

Finally we define the class of proto-quantales and prove some of the remaining facts from the article. Full c-quantales, as defined there, are not needed

for these proofs.

```
class proto-quantale = complete-lattice + proto-monoid +
  assumes Sup-mult-distr: Sup X · y = Sup {x · y | x. x ∈ X}
  and isol: x ≤ y ==> z · x ≤ z · y
```

```
begin
```

```
sublocale pd?: proto-dioid 1σ (·) sup (≤) (<) Sup {}
  ⟨proof⟩
```

```
definition star-rd :: 'a ⇒ 'a where
  star-rd x = Sup {power-rd x i | i. i ∈ ℑ}
```

```
definition star-sq :: 'a ⇒ 'a where
  star-sq x = Sup {power-sq x i | i. i ∈ ℑ}
```

Now we prove Lemma 12.6.

```
lemma star-rd-le-sq: star-rd x ≤ star-sq x
  ⟨proof⟩
```

```
lemma star-sq-le-rd: star-sq x ≤ star-rd x
  ⟨proof⟩
```

```
lemma star-rd-sq: star-rd x = star-sq x
  ⟨proof⟩
```

```
lemma star-sq-power: star-sq x = Sup {pd.p-power (sup 1σ x) i | i. i ∈ ℑ}
  ⟨proof⟩
```

The following lemma should be somewhere close to complete lattices.

```
end
```

```
lemma mono-aux: mono (λy. sup (z: 'a :: proto-quantale) (x · y))
  ⟨proof⟩
```

```
lemma gfp-lfp-prop: sup (gfp (λ(y :: 'a :: proto-quantale). x · y)) (lfp (λy. sup z (x · y))) ≤ gfp (λy. sup z (x · y))
  ⟨proof⟩
```

```
end
```

### 3 Multirelations

```
theory Multirelations
```

```
imports C-Algebras
```

```
begin
```

### 3.1 Basic Definitions

We define a type synonym for multirelations.

**type-synonym**  $'a\ mrel = ('a * ('a set))\ set$

**no-notation**  $s\text{-prod}$  (**infixl**  $\leftrightarrow 80$ )  
**no-notation**  $s\text{-id}$  ( $\langle 1_\sigma \rangle$ )  
**no-notation**  $c\text{-prod}$  (**infixl**  $\langle\parallel\rangle 80$ )  
**no-notation**  $c\text{-id}$  ( $\langle 1_\pi \rangle$ )

Now we start with formalising the multirelational model. First we define sequential composition and parallel composition of multirelations, their units and the universal multirelation as in Section 2 of the article.

**definition**  $s\text{-prod} :: 'a\ mrel \Rightarrow 'a\ mrel \Rightarrow 'a\ mrel$  (**infixl**  $\leftrightarrow 70$ ) **where**  
 $R \cdot S = \{(a,A). (\exists B. (a,B) \in R \wedge (\exists f. (\forall b \in B. (b,f b) \in S) \wedge A = \bigcup \{f b \mid b \in B\}))\}$

**definition**  $s\text{-id} :: 'a\ mrel$  ( $\langle 1_\sigma \rangle$ ) **where**  
 $1_\sigma \equiv \bigcup a. \{(a,\{a\})\}$

**definition**  $p\text{-prod} :: 'a\ mrel \Rightarrow 'a\ mrel \Rightarrow 'a\ mrel$  (**infixl**  $\langle\parallel\rangle 70$ ) **where**  
 $R \parallel S = \{(a,A). (\exists B C. A = B \cup C \wedge (a,B) \in R \wedge (a,C) \in S)\}$

**definition**  $p\text{-id} :: 'a\ mrel$  ( $\langle 1_\pi \rangle$ ) **where**  
 $1_\pi \equiv \bigcup a. \{(a,\{\})\}$

**definition**  $U :: 'a\ mrel$  **where**  
 $U \equiv \{(a,A) \mid a. a \in UNIV \wedge A \subseteq UNIV\}$

**abbreviation**  $NC \equiv U - 1_\pi$

We write NC where  $\overline{1_\pi}$  is written in [2].

Next we prove some basic set-theoretic properties.

**lemma**  $s\text{-prod-im}$ :  $R \cdot S = \{(a,A). (\exists B. (a,B) \in R \wedge (\exists f. (\forall b \in B. (b,f b) \in S) \wedge A = \bigcup ((\lambda x. f x) ` B)))\}$   
 $\langle proof \rangle$

**lemma**  $s\text{-prod-iff}$ :  $(a,A) \in (R \cdot S) \longleftrightarrow (\exists B. (a,B) \in R \wedge (\exists f. (\forall b \in B. (b,f b) \in S) \wedge A = \bigcup ((\lambda x. f x) ` B)))$   
 $\langle proof \rangle$

**lemma**  $s\text{-id-iff}$ :  $(a,A) \in 1_\sigma \longleftrightarrow A = \{a\}$   
 $\langle proof \rangle$

**lemma**  $p\text{-prod-iff}$ :  $(a,A) \in R \parallel S \longleftrightarrow (\exists B C. A = B \cup C \wedge (a,B) \in R \wedge (a,C) \in S)$   
 $\langle proof \rangle$

```

named-theorems mr-simp
declare s-prod-im [mr-simp] p-prod-def [mr-simp] s-id-def [mr-simp] p-id-def [mr-simp]
U-def [mr-simp]

```

## 3.2 Multirelations and Proto-Dioids

We can now show that multirelations form proto-trioids. This is Proposition 5.1, and it subsumes Proposition 4.1,

**interpretation** mrelproto-trioid: proto-trioid  $1_\sigma$   $(\cdot)$   $1_\pi$   $(\parallel)$   $(\cup)$   $(\subseteq)$   $(\subset)$   $\{\}$   
 $\langle proof \rangle$

## 3.3 Simple Properties

This covers all the identities in the display before Lemma 2.1 except the two following ones.

**lemma** s-prod-assoc1:  $(R \cdot S) \cdot T \subseteq R \cdot (S \cdot T)$   
 $\langle proof \rangle$

**lemma** seq-conc-subdistr:  $(R \parallel S) \cdot T \subseteq (R \cdot T) \parallel (S \cdot T)$   
 $\langle proof \rangle$

Next we provide some counterexamples. These do not feature in [2].

**lemma**  $R \cdot \{\} = \{\}$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $R \cdot (S \cup T) = R \cdot S \cup R \cdot T$   
 $\langle proof \rangle$

**lemma**  $R \cdot (S \cdot T) \subseteq (R \cdot S) \cdot T$   
 $\langle proof \rangle$

**lemma**  $(R \parallel R) \cdot T = (R \cdot T) \parallel (R \cdot T)$   
**quickcheck**  
 $\langle proof \rangle$

Next we prove the distributivity and associativity laws for sequential subidentities mentioned before Lemma 2.1

**lemma** subid-aux2:  
**assumes**  $R \subseteq 1_\sigma$  **and**  $(a, A) \in R$   
**shows**  $A = \{a\}$   
 $\langle proof \rangle$

**lemma** s-prod-test-aux1:  
**assumes**  $S \subseteq 1_\sigma$   
**and**  $(a, A) \in R \cdot S$   
**shows**  $((a, A) \in R \wedge (\forall a \in A. (a, \{a\}) \in S))$

```

⟨proof⟩

lemma s-prod-test-aux2:
assumes (a,A) ∈ R
and ∀ a ∈ A. (a,{a}) ∈ S
shows (a,A) ∈ R · S
⟨proof⟩

lemma s-prod-test:
assumes P ⊆ 1σ
shows (a,A) ∈ R · P ←→ (a,A) ∈ R ∧ (∀ a ∈ A. (a,{a}) ∈ P)
⟨proof⟩

lemma test-s-prod-aux1:
assumes P ⊆ 1σ
and (a,A) ∈ P · R
shows (a,{a}) ∈ P ∧ (a,A) ∈ R
⟨proof⟩

lemma test-s-prod-aux2:
assumes (a,A) ∈ R
and (a,{a}) ∈ P
shows (a,A) ∈ P · R
⟨proof⟩

lemma test-s-prod:
assumes P ⊆ 1σ
shows (a,A) ∈ P · R ←→ (a,{a}) ∈ P ∧ (a,A) ∈ R
⟨proof⟩

lemma test-assoc1:
assumes P ⊆ 1σ
shows (R · P) · S = R · (P · S)
⟨proof⟩

lemma test-assoc2:
assumes P ⊆ 1σ
shows (P · R) · S = P · (R · S)
⟨proof⟩

lemma test-assoc3:
assumes P ⊆ 1σ
shows (R · S) · P = R · (S · P)
⟨proof⟩

lemma s-distl-test:
assumes R ⊆ 1σ
shows R · (S ∪ T) = R · S ∪ R · T
⟨proof⟩

```

Next we verify Lemma 2.1.

**lemma** *subid-par-idem*:

**assumes**  $R \subseteq 1_\sigma$

**shows**  $R \parallel R = R$

$\langle proof \rangle$

**lemma** *term-par-idem*:

**assumes**  $R \subseteq 1_\pi$

**shows**  $R \parallel R = R$

$\langle proof \rangle$

**lemma** *U-par-idem*:  $U \parallel U = U$

$\langle proof \rangle$

**lemma** *nc-par-idem*:  $NC \parallel NC = NC$

$\langle proof \rangle$

Next we prove the properties of Lemma 2.2 and 3.2. First we prepare to show that multirelations form c-lattices.

We define the domain operation on multirelations and verify the explicit definition from Section 3.

**definition**  $d :: 'a mrel \Rightarrow 'a mrel$  **where**

$d R \equiv \{(a, \{a\}) \mid a. \exists B. (a, B) \in R\}$

**named-theorems** *mrd-simp*

**declare** *mr-simp* [*mrd-simp*] *d-def* [*mrd-simp*]

**lemma** *d-def-expl*:  $d R = (R \cdot 1_\pi) \parallel 1_\sigma$

$\langle proof \rangle$

**interpretation** *mrel-pbdl-monoid*: *pbdl-monoid*  $1_\sigma$   $(\cdot)$   $1_\pi$   $(\parallel)$   $(\cup)$   $(\subseteq)$   $(\subset)$   $\{\}$   $U$

$(\cap)$

$\langle proof \rangle$

Here come the properties of Lemma 2.2.

**lemma** *c1*:  $(R \cdot 1_\pi) \parallel R = R$

$\langle proof \rangle$

**lemma** *t-aux*:  $T \parallel T \subseteq T \implies (\forall a B C. (a, B) \in T \wedge (a, C) \in T \implies (a, B \cup C)$

$\in T)$

$\langle proof \rangle$

**lemma** *cl4*:

**assumes**  $T \parallel T \subseteq T$

**shows**  $(R \cdot T) \parallel (S \cdot T) \subseteq (R \parallel S) \cdot T$

$\langle proof \rangle$

**lemma** *cl3*:  $R \cdot (S \parallel T) \subseteq (R \cdot S) \parallel (R \cdot T)$   
 $\langle proof \rangle$

**lemma** *cl5*:  $(R \cdot S) \cdot (T \cdot \{\}) = R \cdot (S \cdot (T \cdot \{\}))$   
 $\langle proof \rangle$

We continue verifying other c-lattice axioms

**lemma** *cl8-var*:  $d R \cdot S = (R \cdot 1_\pi) \parallel S$   
 $\langle proof \rangle$

**lemma** *cl9-var*:  $d (R \cap 1_\sigma) = R \cap 1_\sigma$   
 $\langle proof \rangle$

**lemma** *cl10-var*:  $d (R - 1_\pi) = 1_\sigma \cap ((R - 1_\pi) \cdot NC)$   
 $\langle proof \rangle$

### 3.4 Multirelations and C-Lattices

Next we show that multirelations form c-lattices (Proposition 7.3) and prove further facts in this setting.

**interpretation** *mrel-c-lattice*: c-lattice  $1_\sigma$  ( $\cdot$ )  $1_\pi$  ( $\parallel$ ) ( $\cup$ ) ( $\subseteq$ ) ( $\subset$ ) ( $\{\}$ )  $U$  ( $\cap$ )  $NC$   
 $\langle proof \rangle$

The following facts from Lemma 2.2 remain to be shown.

**lemma** *p-id-assoc1*:  $(1_\pi \cdot R) \cdot S = 1_\pi \cdot (R \cdot S)$   
 $\langle proof \rangle$

**lemma** *p-id-assoc2*:  $(R \cdot 1_\pi) \cdot T = R \cdot (1_\pi \cdot T)$   
 $\langle proof \rangle$

**lemma** *seq-conc-subdistrl*:  
**assumes**  $P \subseteq 1_\sigma$   
**shows**  $P \cdot (S \parallel T) = (P \cdot S) \parallel (P \cdot T)$   
 $\langle proof \rangle$

**lemma** *test-s-prod-is-meet* [simp]:  
**assumes**  $R \subseteq 1_\sigma$   
**and**  $S \subseteq 1_\sigma$   
**shows**  $R \cdot S = R \cap S$   
 $\langle proof \rangle$

**lemma** *test-p-prod-is-meet*:  
**assumes**  $R \subseteq 1_\sigma$   
**and**  $S \subseteq 1_\sigma$   
**shows**  $R \parallel S = R \cap S$   
 $\langle proof \rangle$

**lemma** *test-multiplicativer*:

**assumes**  $R \subseteq 1_\sigma$   
**and**  $S \subseteq 1_\sigma$   
**shows**  $(R \cap S) \cdot T = (R \cdot T) \cap (S \cdot T)$   
*⟨proof⟩*

Next we verify the remaining fact from Lemma 2.2; in fact it follows from the corresponding theorem of c-lattices.

**lemma** *c6*:  $R \cdot 1_\pi \subseteq 1_\pi$   
*⟨proof⟩*

Next we verify Lemma 3.1.

**lemma** *p-id-st*:  $R \cdot 1_\pi = \{(a,\{\}) \mid a. \exists B. (a,B) \in R\}$   
*⟨proof⟩*

**lemma** *p-id-zero*:  $R \cap 1_\pi = R \cdot \{\}$   
*⟨proof⟩*

**lemma** *p-id-zero-st*:  $R \cap 1_\pi = \{(a,\{\}) \mid a. (a,\{\}) \in R\}$   
*⟨proof⟩*

**lemma** *s-id-st*:  $R \cap 1_\sigma = \{(a,\{a\}) \mid a. (a,\{a\}) \in R\}$   
*⟨proof⟩*

**lemma** *U-seq-st*:  $(a,A) \in R \cdot U \longleftrightarrow (A = \{\} \wedge (a,\{\}) \in R) \vee (\exists B. B \neq \{\} \wedge (a,B) \in R)$   
*⟨proof⟩*

**lemma** *U-par-st*:  $(a,A) \in R \parallel U \longleftrightarrow (\exists B. B \subseteq A \wedge (a,B) \in R)$   
*⟨proof⟩*

Next we verify the relationships after Lemma 3.1.

**lemma** *s-subid-iff1*:  $R \subseteq 1_\sigma \longleftrightarrow R \cap 1_\sigma = R$   
*⟨proof⟩*

**lemma** *s-subid-iff2*:  $R \subseteq 1_\sigma \longleftrightarrow d R = R$   
*⟨proof⟩*

**lemma** *p-subid-iff*:  $R \subseteq 1_\pi \longleftrightarrow R \cdot 1_\pi = R$   
*⟨proof⟩*

**lemma** *vec-iff1*:  
**assumes**  $\forall a. (\exists A. (a,A) \in R) \longrightarrow (\forall A. (a,A) \in R)$   
**shows**  $(R \cdot 1_\pi) \parallel U = R$   
*⟨proof⟩*

**lemma** *vec-iff2*:  
**assumes**  $(R \cdot 1_\pi) \parallel U = R$   
**shows**  $(\forall a. (\exists A. (a,A) \in R) \longrightarrow (\forall A. (a,A) \in R))$   
*⟨proof⟩*

**lemma** *vec-iff*:  $(\forall a. (\exists A. (a,A) \in R) \longrightarrow (\forall A. (a,A) \in R)) \longleftrightarrow (R \cdot 1_\pi) \parallel U = R$   
 $\langle proof \rangle$

**lemma** *ucl-iff*:  $(\forall a A B. (a,A) \in R \wedge A \subseteq B \longrightarrow (a,B) \in R) \longleftrightarrow R \parallel U = R$   
 $\langle proof \rangle$

**lemma** *nt-iff*:  $R \subseteq NC \longleftrightarrow R \cap NC = R$   
 $\langle proof \rangle$

Next we provide a counterexample for the final paragraph of Section 3.

**lemma**  $1_\sigma \cap R \cdot U = R$   
**nitpick**  
 $\langle proof \rangle$

Next we present a counterexample for vectors mentioned before Lemma 9.3.

**lemma**  $d (d R \cdot U) \cdot (d S \cdot U) \cdot U = (d R \cdot U) \cdot (d S \cdot U)$   
**nitpick**  
 $\langle proof \rangle$

Next we prove Tarski' rule (Lemma 9.3).

**lemma** *tarski-aux*:  
**assumes**  $R - 1_\pi \neq \{\}$   
**and**  $(a,A) \in NC$   
**shows**  $(a,A) \in NC \cdot ((R - 1_\pi) \cdot NC)$   
 $\langle proof \rangle$

**lemma** *tarski*:  
**assumes**  $R - 1_\pi \neq \{\}$   
**shows**  $NC \cdot ((R - 1_\pi) \cdot NC) = NC$   
 $\langle proof \rangle$

Next we verify the assumptions of Proposition 9.8.

**lemma** *d-assoc1*:  $d R \cdot (S \cdot T) = (d R \cdot S) \cdot T$   
 $\langle proof \rangle$

**lemma** *d-meet-distr-var*:  $(d R \cap d S) \cdot T = (d R \cdot T) \cap (d S \cdot T)$   
 $\langle proof \rangle$

Lemma 10.5.

**lemma**  $((R \cap 1_\sigma) \cdot (S \cap 1_\sigma)) \cdot 1_\pi = ((R \cap 1_\sigma) \cdot 1_\pi) \cdot ((S \cap 1_\sigma) \cdot 1_\pi)$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $d ((R \cdot 1_\pi) \cdot (S \cdot 1_\pi)) = d (R \cdot 1_\pi) \cdot d (S \cdot 1_\pi)$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $((R \cap 1_\sigma) \cdot (S \cap 1_\sigma)) \cdot U = ((R \cap 1_\sigma) \cdot U) \cdot ((S \cap 1_\sigma) \cdot U)$

**nitpick**

$\langle proof \rangle$

**lemma**  $d (((R \cdot 1_\pi) \parallel U) \cdot ((S \cdot 1_\pi) \parallel U)) = d ((R \cdot 1_\pi) \parallel U) \cdot d ((S \cdot 1_\pi) \parallel U)$

**nitpick**

$\langle proof \rangle$

**lemma**  $((R \cdot 1_\pi) \cdot (S \cdot 1_\pi)) \parallel U = ((R \cdot 1_\pi) \parallel U) \cdot ((S \cdot 1_\pi) \parallel U)$

**nitpick**

$\langle proof \rangle$

**lemma**  $(((R - 1_\pi) \cap 1_\sigma) \cdot ((S - 1_\pi) \cap 1_\sigma)) \cdot 1_\pi = (((R - 1_\pi) \cap 1_\sigma) \cdot 1_\pi) \cdot (((S - 1_\pi) \cap 1_\sigma) \cdot 1_\pi)$

**nitpick**

$\langle proof \rangle$

**lemma**  $d (((R - 1_\pi) \cdot 1_\pi) \cdot ((S - 1_\pi) \cdot 1_\pi)) = d ((R - 1_\pi) \cdot 1_\pi) \cdot d ((S - 1_\pi)$

$\cdot 1_\pi)$

**nitpick**

$\langle proof \rangle$

**lemma**  $(((R - 1_\pi) \cap 1_\sigma) \cdot ((S - 1_\pi) \cap 1_\sigma)) \cdot NC = (((R - 1_\pi) \cap 1_\sigma) \cdot NC) \cdot (((S - 1_\pi) \cap 1_\sigma) \cdot NC)$

$\langle proof \rangle$

**lemma**  $d (((((R - 1_\pi) \cdot 1_\pi) \parallel NC) \cdot (((S - 1_\pi) \cdot 1_\pi) \parallel NC)) = d (((R - 1_\pi) \cdot$

$1_\pi) \parallel NC) \cdot d (((S - 1_\pi) \cdot 1_\pi) \parallel NC)$

$\langle proof \rangle$

**lemma**  $(((R - 1_\pi) \cdot 1_\pi) \cdot ((S - 1_\pi) \cdot 1_\pi)) \parallel NC = (((R - 1_\pi) \cdot 1_\pi) \parallel NC) \cdot (((S - 1_\pi) \cdot 1_\pi) \parallel NC)$

**nitpick**

$\langle proof \rangle$

**lemma**  $(((R - 1_\pi) \cdot 1_\pi) \parallel NC) \cdot (((S - 1_\pi) \cdot 1_\pi) \parallel NC) \cdot 1_\pi = (((((R - 1_\pi) \cdot$

$1_\pi) \parallel NC) \cdot 1_\pi) \cdot (((((S - 1_\pi) \cdot 1_\pi) \parallel NC) \cdot 1_\pi)$

**nitpick**

$\langle proof \rangle$

### 3.5 Terminal and Nonterminal Elements

Lemma 11.4

**lemma**  $(R \cdot S) \cdot \{\} = (R \cdot \{\}) \cdot (S \cdot \{\})$

**nitpick**

$\langle proof \rangle$

**lemma**  $(R \cdot S) - 1_\pi = (R - 1_\pi) \cdot (S - 1_\pi)$

$\langle proof \rangle$

**lemma**  $(R \parallel S) - 1_\pi = (R - 1_\pi) \parallel (S - 1_\pi)$   
**nitpick**  
 $\langle proof \rangle$

Lemma 11.8.

**lemma**  $((R \cdot 1_\pi) \cdot (S - 1_\pi)) - 1_\pi = (R \cdot 1_\pi) \cdot (S - 1_\pi)$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $((S - 1_\pi) \cdot (R \cdot 1_\pi)) - 1_\pi = (S - 1_\pi) \cdot (R \cdot 1_\pi)$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $((R \cdot 1_\pi) \parallel (S - 1_\pi)) \cdot 1_\pi = (R \cdot 1_\pi) \parallel (S - 1_\pi)$   
**nitpick**  
 $\langle proof \rangle$

Lemma 11.10.

**lemma**  $R \cdot \{\} \subseteq S \cdot \{\} \implies (R \cdot T) \cdot \{\} \subseteq (S \cdot T) \cdot \{\}$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $R - 1_\pi \subseteq S - 1_\pi \implies (R \parallel T) - 1_\pi \subseteq (S \parallel T) - 1_\pi$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $R - 1_\pi \subseteq S - 1_\pi \implies (T \cdot R) - 1_\pi \subseteq (T \cdot S) - 1_\pi$   
 $\langle proof \rangle$

Corollary 11.12

**lemma**  $R \cdot \{\} = S \cdot \{\} \implies (R \cdot T) \cdot \{\} = (S \cdot T) \cdot \{\}$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $R - 1_\pi = S - 1_\pi \implies (R \parallel T) - 1_\pi = (S \parallel T) - 1_\pi$   
**nitpick**  
 $\langle proof \rangle$

**lemma**  $R - 1_\pi = S - 1_\pi \implies (T \cdot R) - 1_\pi = (T \cdot S) - 1_\pi$   
 $\langle proof \rangle$

### 3.6 Multirelations, Proto-Quantales and Iteration

**interpretation** *mrel-proto-quantale*: proto-quantale  $1_\sigma$   $(\cdot)$  Inter Union  $(\cap)$   $(\subseteq)$   
 $(\subset)$   $(\cup)$   $\{\}$   $U$   
 $\langle proof \rangle$

We reprove Corollary 13.2. because Isabelle does not pick it up from the quantale level.

**lemma** *iso-prop: mono* ( $\lambda X. S \cup R \cdot X$ )  
*(proof)*

**lemma** *gfp-lfp-prop: gfp* ( $\lambda X. R \cdot X$ )  $\cup$  *lfp* ( $\lambda X. S \cup R \cdot X$ )  $\subseteq$  *gfp* ( $\lambda X. S \cup R \cdot X$ )  
*(proof)*

### 3.7 Further Counterexamples

Lemma 14.1. and 14.2

**lemma**  $R \parallel R \subseteq R$   
**nitpick**  
*(proof)*

**lemma**  $R \subseteq R \parallel S$   
**nitpick**  
*(proof)*

**lemma**  $R \parallel S \cap R \parallel T \subseteq R \parallel (S \cap T)$   
**nitpick**  
*(proof)*

**lemma**  $R \cdot (S \parallel T) = (R \cdot S) \parallel (R \cdot T)$   
**nitpick**  
*(proof)*

**lemma**  $R \cdot (S \cdot T) \subseteq (R \cdot S) \cdot T$   
*(proof)*

**lemma**  $\llbracket R \parallel R = R; S \parallel S = S; T \parallel T = T \rrbracket \implies R \cdot (S \parallel T) = (R \cdot S) \parallel (R \cdot T)$   
**nitpick**  
*(proof)*

**lemma**  $\llbracket R \neq \{\}; S \neq \{\}; \forall a. (a, \{\}) \notin R \cup S \rrbracket \implies R \cdot S \subseteq R \parallel S$   
**quickcheck**  
*(proof)*

**lemma**  $\llbracket R \neq \{\}; S \neq \{\}; \forall a. (a, \{\}) \notin R \cup S \rrbracket \implies R \parallel S \subseteq R \cdot S$   
**quickcheck**  
*(proof)*

**lemma**  $\llbracket R \neq \{\}; S \neq \{\}; T \neq \{\}; \forall a. (a, \{\}) \notin R \cup S \cup T \rrbracket \implies (R \parallel S) \cdot T \subseteq R \parallel (S \cdot T)$   
**quickcheck**  
*(proof)*

```

lemma  $\llbracket R \neq \{\}; S \neq \{\}; T \neq \{\}; \forall a. (a, \{\}) \notin R \cup S \cup T \rrbracket \implies R \parallel (S \cdot T) \subseteq (R \parallel S) \cdot T$ 
quickcheck
 $\langle proof \rangle$ 

lemma  $\llbracket R \neq \{\}; S \neq \{\}; T \neq \{\}; \forall a. (a, \{\}) \notin R \cup S \cup T \rrbracket \implies R \cdot (S \parallel T) \subseteq (R \cdot S) \parallel T$ 
quickcheck
 $\langle proof \rangle$ 

lemma  $\llbracket R \neq \{\}; S \neq \{\}; T \neq \{\}; \forall a. (a, \{\}) \notin R \cup S \cup T \rrbracket \implies (R \cdot S) \parallel T \subseteq R \cdot (S \parallel T)$ 
quickcheck
 $\langle proof \rangle$ 

lemma  $\llbracket R \neq \{\}; S \neq \{\}; \forall a. (a, \{\}) \notin R \cup S \rrbracket \implies (R \parallel S) \cdot (R \parallel S) \subseteq (R \cdot R) \parallel (S \cdot S)$ 
quickcheck
 $\langle proof \rangle$ 

```

```

lemma  $\llbracket R \neq \{\}; S \neq \{\}; \forall a. (a, \{\}) \notin R \cup S \rrbracket \implies (R \cdot R) \parallel (S \cdot S) \subseteq (R \parallel S) \cdot (R \parallel S)$ 
quickcheck
 $\langle proof \rangle$ 

```

### 3.8 Relationship with Up-Closed Multirelations

We now define Parikh's sequential composition.

```

definition s-prod-pa :: 'a mrel  $\Rightarrow$  'a mrel  $\Rightarrow$  'a mrel (infixl  $\langle\otimes\rangle$  70) where
   $R \otimes S = \{(a, A). (\exists B. (a, B) \in R \wedge (\forall b \in B. (b, A) \in S))\}$ 

```

We show that Parikh's definition doesn't preserve up-closure.

```

lemma up-closed-prop:  $((R \parallel U) \cdot (S \parallel U)) \parallel U = (R \parallel U) \cdot (S \parallel U)$ 
 $\langle proof \rangle$ 

```

Lemma 15.1.

```

lemma onelem:  $(R \cdot S) \parallel U \subseteq R \otimes (S \parallel U)$ 
 $\langle proof \rangle$ 

```

```

lemma twolem:  $R \otimes (S \parallel U) \subseteq (R \cdot S) \parallel U$ 
 $\langle proof \rangle$ 

```

```

lemma pe-pa-sim:  $(R \cdot S) \parallel U = R \otimes (S \parallel U)$ 
 $\langle proof \rangle$ 

```

```

lemma pe-pa-sim-var:  $((R \parallel U) \cdot (S \parallel U)) \parallel U = (R \parallel U) \otimes (S \parallel U)$ 
 $\langle proof \rangle$ 

```

**lemma** *pa-assoc1*:  $((R \parallel U) \otimes (S \parallel U)) \otimes (T \parallel U) \subseteq (R \parallel U) \otimes ((S \parallel U) \otimes (T \parallel U))$   
 $\langle proof \rangle$

The converse direction of associativity remains to be proved.

Corollary 15.3.

**lemma** *up-closed-par-is-meet*:  $(R \parallel U) \parallel (S \parallel U) = (R \parallel U) \cap (S \parallel U)$   
 $\langle proof \rangle$

**end**

## References

- [1] H. Furusawa and G. Struth. Concurrent dynamic algebra. *ACM Transactions on Computational Logic*, 2015. (In Press).
- [2] H. Furusawa and G. Struth. Taming multirelations. *CoRR*, abs/1501.05147, 2015.
- [3] D. Peleg. Concurrent dynamic logic. *J. ACM*, 34(2):450–479, 1987.